

Chroniberia: The Ongoing Development of a Geochronological GIS Database of Iberia

C. Lopes, A. Ferreira, M. Chichorro, M. F. Pereira, J. A. Almeida and A. R Solá

Abstract Geochronological comparisons of large datasets are facilitated by the use of structured databases. This paper reports the ongoing development of CHRONIBERIA, a geochronological GIS database that will provide access to zircon U–Pb age datasets from Iberia. The database will also integrate relevant distinct datasets from other correlative regions around the world to construct a framework based on stratigraphic, sedimentological, and palaeogeographical data.

Keywords Geochronology · Zircon U–Th–Pb · Database · GIS · Iberia

A detailed and precise temporal framework is essential for exploring the richness of the geological record (Bowring and Schmitz 2003). Zircon U–Th–Pb geochronology has been widely used to provide an exact measure of time (Davis et al. 2003), which has give rise to large geochronological datasets. Geochronological comparisons of such large datasets are facilitated by the use of structured databases. Several distinct geochronological databases and database systems have been developed in the last decade, including: CHRONOBANK, a GIS-based geochronological database (Brazil, Silva et al. 2003); NGDB, the National Geochronological Database (USA, Silva et al. 2003); and OZCHRON (Australia, Sircombe 2006). DateView (a freeware desktop database system; Eglington 2004), the StratDB/IGCP509 initiative (a database system and web interface with links to the geochronology database DateView; Eglington et al. 2009), and the EarthChem initiative (<http://www.earthchem.org/earthchemWeb/index.jsp>) are all part of the

C. Lopes · A. Ferreira · M. Chichorro (✉) · J. A. Almeida
CICEGe, Departamento de Ciências da Terra, Faculdade de Ciências e Tecnologia,
Universidade Nova de Lisboa, 2829-516 Caparica, Portugal
e-mail: ma.chichorro@fct.unl.pt

M. F. Pereira
IDL, Departamento de Geociências, ECT, Universidade de Évora, Évora, Portugal

A. R Solá
Unidade de Geologia, Hidrogeologia e Geologia Costeira, LNEG, Lisbon, Portugal

most recent developments to construct an international database system. These databases use GIS tools to improve spatial analysis, to generate maps, and to integrate distinct data set (Silva et al. 2003; Eglinton 2004). To understand the geological history of the Gondwana supercontinent and related peripheral magmatic arcs (the Avalonia and Cadomian terranes), a geochronological GIS database of Iberia (CHRONIBERIA), based on the zircon contents of rocks from Iberia, has been created.

Design of the Geochronological Database

In the conception of the database, the following main issues were considered: the nature of the data; the types of queries expected; and the workflow and main goals of the GONDWANA project (a research project that focuses on obtaining new U–Pb zircon ages in southwestern Iberia and which intends to analyse them together with data already published on Iberia). Some technical considerations were also addressed: the number of users and usage pattern; the type of database (relational); the usage of the database (for editing/read-only); the degree of demand (high/low); the type of software; the connection/transaction to the web; and the administration of the database.

Based on a set of typical queries normally applied to isotope data (see Eglinton 2004), the nature, detail, and amount of information to be used in the CHRONIBERIA were, on a first level, defined as follows: Which samples are associated with some specified age? Which samples from a specific formation are associated with some specified age? What is the age of a specified geological unit or of an event? How do dates from one area or geological entity compare with those from other areas? What is the geographical distribution of dates in some area relative to their geological interpretation? How do initial ratios vary with time in some areas? It is clear that spatial information is required in addition to chronological.

Based on a synthetic workflow related to isotope data acquisition, the major entities and relations of the database were defined (Fig. 1). The schematic design of the database was subsequently defined based on a diagram of entities and relationships.

Some technical considerations were then addressed, regarding the type of software. Databases using Microsoft Access offer, for example, better and more user-friendly manipulation of alphanumeric data and easier generation of data forms and queries involving several tables, whereas ArcGIS databases (geodatabases) provide the possibility of making spatial queries of and spatial operations on the data. Original datafile formats for ArcGIS geodatabases can be stored and managed in Microsoft Access data files, which offers a hybrid solution as a starting point.

developed and tested. Moreover, a WebGIS (Geographic Information System for the World Wide Web) will be implemented.

Acknowledgments C. Lopes acknowledges a research grant of 12 months duration that was sponsored by project GONDWANA-PTDC/CTE-GIX/110426/2009 funded by Fundação para a Ciência e Tecnologia (Portugal). It is a contribution to project PESt-OE/CTE/UI4073/2014.

References

- Bowring, S. A., & Schmitz, M. D. (2003). High-precision U-Pb zircon geochronology and the stratigraphic record. *Reviews in Mineralogy and Geochemistry*, 53, 305–326.
- Chichorro, M., Pereira, M. F., Díaz-Azpiroz, M., Williams, I. S., Fernández, C., Pin, C., et al. (2008). Cambrian ensialic rift-related magmatism in the Ossa-Morena Zone (Évora–Aracena metamorphic belt, SW Iberian Massif): Sm–Nd isotopes and SHRIMP zircon U–Th–Pb geochronology. *Tectonophysics*, 461, 91–113.
- Davis, D., Williams, I. S., & Krogh, T. E. (2003). Historical development of zircon geochronology. *Reviews in Mineralogy and Geochemistry*, 53, 145–182.
- Eglington, B. M. (2004). DateView: a windows geochronology database. *Computers & Geosciences*, 30, 847–858.
- Eglington, B. M., Reddy, S. M., & Evans, D. A. D. (2009). The IGCP 509 database system; design and application of a tool to capture and illustrate litho- and chrono-stratigraphic information for Palaeoproterozoic tectonic domains, large igneous provinces and ore deposits; with examples from Southern Africa (in Palaeoproterozoic supercontinents and global evolution). *Geological Society Special Publications*, 323, 27–47.
- Pereira, M. F., Solá, A. R., Chichorro, M., Lopes, L., Gerde, A., & Silva, J. B. (2012). North-Gondwana assembly, break up and paleogeography: U–Pb isotope evidence from detrital and igneous zircons of Ediacaran and Cambrian rocks of SW Iberia. *Gondwana Research* 22(3–4), 866–881. <http://dx.doi.org/10.1016/j.gr.2012.02.010>
- Silva, L. C., Rodrigues, J. B., Silveira, L. M. C., & Pimentel, M. M. (2003). The Brazilian National Geochronological Database: Chronobank, *IV South American Symposium on Isotope Geology*, Salvador.
- Sircombe, K. N. (2006). Standardising geochronological data. *Geochimica et Cosmochimica Acta*, 70, A594.
- Solá, A. R., Pereira, M. F., Williams, I. S., Ribeiro, M. L., Neiva, A. M. R., Montero, P., et al. (2008). New insights from U–Pb zircon dating of Early Ordovician magmatism on the northern Gondwana margin: The Urro Formation (SW Iberian Massif, Portugal). *Tectonophysics*, 461, 114–129.