



## OPEN ACCESS

## EDITED BY

Rachid Bouharroud,  
National Institute of Agronomic Research,  
Morocco

## REVIEWED BY

Mohammed Antar,  
McGill University, Macdonald Campus,  
Canada  
Luciane Maria Colla,  
The University of Passo Fundo, Brazil

## \*CORRESPONDENCE

Alice Ferreira

✉ [alice.ferreira@lneg.pt](mailto:alice.ferreira@lneg.pt)

<sup>†</sup>These authors have contributed equally to  
this work

RECEIVED 07 October 2022

ACCEPTED 25 May 2023

PUBLISHED 09 June 2023

## CITATION

Ferreira A, Bastos CRV,  
Marques-dos-Santos C,  
Acién-Fernandez FG and  
Gouveia L (2023) Algaeculture for  
agriculture: from past to future.  
*Front. Agron.* 5:1064041.  
doi: 10.3389/fagro.2023.1064041

## COPYRIGHT

© 2023 Ferreira, Bastos,  
Marques-dos-Santos, Acién-Fernandez  
and Gouveia. This is an open-access article  
distributed under the terms of the [Creative  
Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The  
use, distribution or reproduction in other  
forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in  
this journal is cited, in accordance with  
accepted academic practice. No use,  
distribution or reproduction is permitted  
which does not comply with these terms.

# Algaeculture for agriculture: from past to future


Alice Ferreira<sup>1\*†</sup>, Carolina R. V. Bastos<sup>2†</sup>,  
Cláudia Marques-dos-Santos<sup>3</sup>,  
Francisco Gabriel Acién-Fernandez<sup>4</sup> and Luisa Gouveia<sup>1,2</sup>

<sup>1</sup>Bioenergy Unit, LNEG, National Laboratory of Energy and Geology I.P., Lisboa, Portugal, <sup>2</sup>Green  
CoLab, Green Ocean Technologies and Products Collaborative Laboratory, Centro de Ciências do  
Mar do Algarve, Universidade do Algarve, Faro, Portugal, <sup>3</sup>Forest Research Centre, School of  
Agriculture, University of Lisbon, Lisboa, Portugal, <sup>4</sup>Chemical Engineering Department, University of  
Almería, Almería, Spain

The continuous growth of the world population has imposed major challenges on agriculture. Consequently, farmers generalized the overuse of synthetic fertilizers and pesticides to meet the global food demand. Although these products have helped many developing countries increase their crop yield, they have simultaneously resulted in many issues, mainly the decline of soil fertility and degradation of local ecosystems due to soil, water, and air contamination, combined with their non-renewable nature and increased costs. For agriculture to become more sustainable, the use of alternative biological products, with recognized beneficial effects on plant yield and health, must be expanded. In this context, microalgae and cyanobacteria are rich sources of nutrients and bioactive metabolites, which have been gaining attention from researchers and companies for their ability to improve plant nutrition, growth, and tolerance to stress. This review gives an overview of the research work that has been done in the last two decades, regarding the use of microalgae and cyanobacteria (blue-green algae) as biofertilizers, biostimulants, and biopesticides. This work identified trends and challenges and highlights the use of microalgae to recycle the nutrients from wastewater to improve plant productivity while reducing the fertilizer and water footprint for more sustainable agriculture practices.

## KEYWORDS

wastewater, microalgae, cyanobacteria, biostimulants, biofertilizers,  
biopesticides, biocontrol



## Introduction

Agriculture is facing significant pressure to meet the growing food demands of the increasing world population. Until the 20th century, nitrogen and phosphorus were the limiting factors for agricultural yield. Despite the abundance of atmospheric nitrogen, most crop plants cannot uptake it in this form and therefore need the application of fertilizers. Conventional fertilizers are obtained primarily from declining nonrenewable resources. Currently, more than 96% of ammonia is produced through the Haber-Bosch process, which requires fossil fuels as feedstock (natural gas, oil, and coal) (Smith et al., 2020). This process is one of the largest consumers of global energy, expending more than 1–2% of the annual world energy supply (Chatterjee et al., 2017). Furthermore, this process is also a large greenhouse gas emitter, responsible for 1.2% of global anthropogenic CO<sub>2</sub> emissions (Smith et al., 2020).

Regarding phosphorus, all the produced fertilizers are derived from limited mineral deposits which are restricted to specific areas of the planet. Approximately 75% of the current reserves of phosphate minerals are in Morocco. China and the US also have significant reserves, but these are not sold on the global market. This problem of scarcity and the key role of phosphorus in agriculture makes it highly susceptible to unpredictable price fluctuations on the global market, which is also strongly influenced by geopolitical interests. Countries that do not have their reserves, including the European Union are more at risk (Zilio et al., 2022). These concerns are more intensified with the recent war triggered by the Russian invasion of Ukraine.

The productivity of crops has been negatively affected by the appearance of pests and diseases, and increase of water scarcity, due to the soil erosion and degradation of local ecosystems by the intensive use of synthetic fertilizers (Chatterjee et al., 2017). Despite the detrimental environmental impacts, the use of fertilizers is inevitable to meet the growing global food demand. As most environmental legislations worldwide become more rigorous, the emergence of organic farming practices appears as a promising solution to fulfill the growing demand for healthy foods that do not compromise long-term environmental sustainability, promote low pesticide inputs, and the use of natural bioproducts, such as biofertilizers, biostimulants, and biopesticides (Calvo et al., 2014; Bulgari et al., 2015; Colla and Rouphael, 2015; Chatterjee et al., 2017; Win et al., 2018). Still, when it comes to bioproducts, misinterpretations between these concepts can still occur, especially between biofertilizers and biostimulants (Figure 1). According to the new EU fertilizing products regulation (FPR) 2019/1009 that lays down rules on the making available on the market EU fertilising products, and was implemented on 16<sup>th</sup> of July of 2022, a biostimulant is “an EU fertilizing product whose function is to stimulate plant nutrition processes independently of the product’s nutrient content with the sole aim of improving one or more of the following characteristics of the plant or the plant rhizosphere: nutrient use efficiency, tolerance to abiotic stress, quality traits, or availability of confined nutrients in the soil or rhizosphere” (Regulation (EU) 2019/1009, 2019). Essentially, the main difference between both concepts is that biostimulants do not provide nutrients directly to crops as opposed to biofertilizers.




FIGURE 1

Summary of natural bioproducts (biofertilizer, biostimulant, and biocontrol) and their impact on the crop.

Biofertilizers are biological products that contain microorganisms, such as algae, fungi, or bacteria, or natural compounds derived from those microorganisms, which act at the soil level. These products promote plant growth and soil fertility by increasing the supply or availability of essential nutrients, including macro and micronutrients, and by improving physico-chemical and biological soil properties (Abdel-Raouf, 2012; Sahu et al., 2012; Ronga et al., 2019). On the other hand, biostimulants stimulate plant growth by modulating plant growth metabolisms, such as germination, respiration, photosynthetic activity, nutrient uptake, and flower and fruit production, amongst others (Górka et al., 2018). They also play a relevant role in extending plant tolerance and resistance to various environmental stress conditions (e.g., drought, heat, salinity, etc.) (Povero et al., 2016; Chiaiese et al., 2018). On the other hand, biopesticides promote antagonism and biological control of pathogenic organisms, such as bacteria, viruses, fungi, and insects (Carvajal-Muñoz and Carmona-García, 2012).

One group of microorganisms that can act as bioproducts in Agriculture are microalgae. These organisms excrete organic acids, increase phosphorus availability and uptake, provide nitrogen fixation, and enhance the soil content of organic matter to build soil fertility (Górka et al., 2015; Coppens et al., 2016; Khan et al., 2019). Microalgae produce biologically active extracellular substances that influence plant growth, also known as Plant Growth Regulators (PGRs) which act at very low concentrations (i.e., phytohormones, amino acids, polypeptides, polysaccharides, and vitamins), whilst specific eukaryotic microalgae have also shown antibacterial and antifungal activity. (Chiaiese et al., 2018; Chanda et al., 2019).

Several bioactive compounds are behind these antimicrobial mechanisms, such as fatty acids, and phenolic compounds (Renuka et al., 2018; Costa et al., 2019). Cyanobacteria have also been reported as stimulators of plant defense mechanisms and producers of metabolites with pesticidal action (Singh et al., 2016). Further research is required for green microalgae on the control of plant pathogens. Nevertheless, studies already reveal that green microalgae (e.g., *Coccomyxa onubensis*, *Chlorella vulgaris*, etc.) have antimicrobial properties, which makes them an interesting subject for research in the agricultural biotechnology field (Costa et al., 2019).

From the economic point of view, the growth of the European biostimulants sector has been driven by the increasing importance of organic and sustainable farming as well as the need for enhanced yields (Mordor Intelligence, 2020). Europe is currently the biggest market for biostimulants, with around 8.5 million acres of area treated in 2016 (Liebig et al., 2020). This has increased the need for harmonized European Regulation, concerning the placement of biostimulants in the market. Thus, the new Fertilizing Products Regulation (FPR) (EU) 2019/1009 includes biostimulants as CE-marked fertilizing products meaning that the producer can now officially evaluate and prove that these products meet EU safety, health, or environmental requirements, therefore receiving the official EU conformity certification to be sold in the markets belonging to the European Economic Area (CE stands for “Conformité Européenne,” which translates to “European Conformity.”) (Regulation (EU) 2019/1009, 2019). The Global Biostimulant Market was estimated to be valued at USD 2.6 billion, in 2019, and is expected to grow by 11.24% by 2025 (MarketsandMarkets, 2020). A recent event that has further

aggravated the expectations and potential of the biostimulant market is the Russian invasion of Ukraine. The beginning of the war in March of 2022, has led to a shortage of fertilizers in the market. Both countries were reported to be responsible, altogether, for the exportation of 28% of fertilizers containing nitrogen, phosphorus, and potassium, according to Morgan Stanley (Domm, 2022). This has resulted in a significant increase in fertilizer prices, which have become worryingly scarce. For the farmers, these events translate to lower crop yields, and for the consumers, increased market prices of the final products. Although the duration of the conflict cannot be foretold with complete certainty, the negative effects on the global markets are expected to last for the coming years, with or without the prolongation of the war. Furthermore, a reshape of the global order is already at hand, with several world leaders working on decreasing the dependency on Russia's supplies, such as fertilizers. This conflict has thus begun a new cycle in the world's economy and the search for alternative suppliers is at its peak. In the case of fertilizers, the current crisis is further aggravated by the climate crisis as the chemicals associated with fertilizers, as well as the production of fertilizers, are highly pollutant. Hence, microalgae bioproducts such as biostimulants and biofertilizers may very well pose an opportunity to alleviate and overcome the current fertilizer crisis.

This manuscript gives an overview of the trends of microalgae, eukaryotic, and cyanobacteria, for agricultural applications, through a literature review of the most relevant studies in this field. The authors pretend to emphasize the relevance of microalgae biomass as an emerging application in agriculture, especially after the use of microalgae for the treatment of wastewater from various sources.

## Database

The data was collected from the Web of Science database. The search was done using several combinations of keywords: (\*microalga\* OR \*cyanobacteri\*) AND (\*biostimulant\* OR \*biofertilizer\* OR \*biopesticide\* OR “germination” OR “plant growth” OR “bioprotection” OR “agriculture”). The search generated 1873 articles from the Science Citation Index (SCI) starting in 2001 until September 2022. From those articles, both reviews and viewpoint articles were removed, and only English language journals were considered (1458 articles). From these articles, we searched within the references for more articles that were not in our initial dataset. Among these, a selection was made to consider only articles which studied the effect of microalgae and cyanobacteria on target plants, soil, or pests. The final dataset included 330 original research papers for a 21-year period (2001 – 2022), which were synthesized in the sections below. Furthermore, an online page displaying the information within this manuscript was prepared at the following link: <https://shorturl.at/klxD0>.

## Geographical distribution of publications

In the last 5 years, the research on the agricultural application of microalgae has registered a steep growth, duplicating the number of

studies in 2021 compared to 2018 (Figure 2A). This demonstrates the emergence of microalgae in the field of agriculture, which might develop to be one of the most relevant applications in microalgae technology in the following years (Figure 2B).

Asia has led the research in this field, being the source of almost half the studies published between 2001 and 2022, with India standing out among them (87 publications). In the last decade, India has emerged as a major agricultural exporter, ranking first in the world with the highest net cropped area (USGS, 2021). However, the Indian agricultural sector has faced various challenges such as low farm yields, limited water availability, and environmental and soil degradation, amongst others (Chew and Soccio, 2016; Narain, 2020). These could be the major incentives behind extensive research on sustainable ways to overcome these challenges. In Europe, Spain has been the leading country in publications (18 publications), closely followed by Italy (17 publications). In Africa and America, Egypt and Brazil stood out, respectively, with 33 and 26 published studies each (Figure 2C). Spain and Italy have been valuable players in the European agriculture sector, however, Germany and France, represent 51% of the total European Union (EU) utilized agricultural area and 49% of the total EU arable land. Moreover, Italy cut down 20% of the use of pesticides between 2011 and 2018 by substituting them for more sustainable alternatives such as practices like crop rotation and technologies like precision farming making it the most sustainable agriculture sector in Europe (Eurostat, 2020).

## Distribution of microalgae and cyanobacteria in agriculture publications

There has been a predominance of studies addressing cyanobacteria species for agricultural purposes over microalgae (185 vs. 108 studies) (Figure 3A). 35 studies used both microalgae and cyanobacteria species, where 6 took advantage of mixed consortia. Among the microalgal species studied (Figure 3B), there was a clear predominance of the genus *Chlorella* (108), being that almost half of the studies used *Chlorella vulgaris*, followed by *Scenedesmus* (48) (including *Tetrademus* and *Desmodesmus* due to changes in taxonomy). Other microalgal genera have been studied, although to a much lesser extent, such as *Dunaliella*, *Nannochloropsis*, *Porphyridium*, and *Tetraselmis*, among others. Regarding cyanobacteria, *Anabaena* and *Nostoc* were dominant genera, especially in studies dealing with soil-borne cyanobacteria in rice paddy fields in India. In fact, throughout the years, many studies regarding the microbial communities of soils, included both genera as the most found species, due to their ability to fix atmospheric nitrogen. In addition, their significant effect on plant growth can be attributed to the ability of *Anabaena* and *Nostoc* strains to produce growth regulators (e.g. phytohormones like indole-acetic acid) and other secondary metabolites, linked to the biocontrol of diseases, as well as improving soil aggregation through secretion of mucilage and polysaccharides (Prasanna et al., 2011). *Arthrospira* has also been the dominant cyanobacterial genus in most recent studies




FIGURE 2

(A) Evolution of publications on the application of microalgae in agriculture worldwide; (B) Evolution of publications on the various applications of microalgae worldwide; and (C) Number of publications on the application of microalgae in agriculture by country, over 21 years (2001–2022).

(58), especially *Arthrospira platensis* (46). Moreover, in the literature, the higher abundance of cyanobacteria compared to microalgae was related to their identification within various consortia collected from soils, not to their isolated effect. The fact that *Chlorella vulgaris* and *Arthrospira platensis* were the most studied microalga and cyanobacteria species, respectively, was expected given their public acceptance and dominance in the global production market.

Regarding the cultivation medium, most studies used synthetic media to produce the microalgal/cyanobacterial biomass (285), while only 40 publications used biomass that was produced in wastewater (Figure 3C). In terms of composition, these wastewaters are typically richer in nutrients such as nitrogen, phosphate, and

potassium, as well as contain significant contents of solids, dissolved and particular matter, and microorganisms, when compared with synthetic media. It is also interesting to notice that most biomass produced from wastewater (WW) were microalgae, either in isolates or in consortium with other microalgal or cyanobacterial species. From the literature, only 3 studies used cyanobacterial isolates or consortia (Wuang et al., 2016; Rashad et al., 2019; Ferreira et al., 2021), while the remaining used consortia with both microalgae and cyanobacteria, dominated by the former (Mulbry et al., 2005; Mukherjee et al., 2016; Renuka et al., 2016; Castro et al., 2017; Ahn et al., 2020; Supraja et al., 2020). From these microalgae, the most common species that have been found in






FIGURE 3

Distribution in agriculture application studies by (A) type of microorganism; (B) number and genus of the microalgae and cyanobacteria used in the studies; (C) cultivation medium used to grow the biomass.

wastewater treatment plants are *Chlorella* and *Scenedesmus*/ *Tetrademus*, two highly robust and resilient strains. This has resulted in the generalized use of these species in studies involving wastewaters, either in the form of isolates or due to their domination of the cultures. (Ferreira et al., 2019; Navarro-López et al., 2020a; Ferreira et al., 2021; Ranglová et al., 2021; Viegas et al., 2021a; Viegas et al., 2021b; Viegas et al., 2021c). The potential of wastewater-grown microalgae for agriculture was further addressed in section 4.1.

## Agricultural applications of microalgae and cyanobacteria

Microalgae can be used either as pure extracts or as crude algal compost to enhance seed germination, and crop productivity at different levels, by improving soil quality, stimulating growth, and/or expanding protection against stress conditions. Figure 4A shows

that most studies used microalgae/cyanobacteria as a biostimulant (162) rather than a biofertilizer (145). There seems to be some incorrect denomination of the effect due to the misconception between both concepts, as previously mentioned, being that some studies did not distinguish between them and considered only a biofertilizer effect. However, for of Figure 4A, a distinction between them was considered depending on their described effect in the respective studies. In comparison, the biocontrol effect has been less studied, with only 26 studies addressing it. Furthermore, being an emerging topic, most studies have been developed at the laboratory scale (154), especially in plant germination and early development stages. Nonetheless, there are a considerable number of studies performed in pots, both in a greenhouse (104) or outdoors (20), and in field trials (70) (Figure 4B).

The most common mode of application was *via* root (207) (Figure 4C). All studies that applied seed soaking (92) were performed at the laboratory scale. This can be explained by the fact that this method is mainly used for germination experiments,




FIGURE 4

Distribution of studies by (A) the agricultural purpose of the biomass; (B) the experimental scale of the trials; (C) the application mode of the biomass and (D) the growth conditions of the plant trials (stress or no stress).

which are usually the fastest ones to perform when the goal is to first study the biostimulant/biofertilizer potential of a certain compound and microorganism. From the literature, only 2 studies were found regarding the biostimulant (Bayona-Morcillo et al., 2020) and biopesticide (Kim et al., 2020) effects of microalgae when applied *via* foliar and root in greenhouse trials. When comparing foliar spray to soil application, the former provides faster nutrient utilization and correction of nutrient deficiencies, with improvements of N, P and K contents (Dineshkumar et al., 2020a). The number of studies evaluating the potential utilization of microalgae and microalgae-derived compounds, such as polysaccharides and phytohormones, for foliar application was limited (45). This strategy is relatively new and is one of the most innovative agricultural practices, as it is environmentally safe and promotes agricultural sustainability (Ronga et al., 2019). Microalgae-derived extracts, even at low concentrations, can induce an array of physiological plant responses. Since the amount of natural substances in algae is relatively smaller than in mineral fertilizers, the foliar application could be a more suitable method, since plant responses to nutrients supplied *via* foliar sprays are normally more rapid than when applied in soil (Ronga et al., 2019).

Regarding growth conditions (Figure 4D), most studies were developed under controlled environments with optimal growth conditions (262). Still, 40 studies have already tested the potential of microalgae/cyanobacteria for improving plant growth under abiotic stress conditions, especially salinity (20), chemical contamination (9), drought (5), and heat (4). The studies considering biotic stress were related to the biopesticide effect (26).

Concerning the type of crop used in these studies, cereal plants were the most common, especially rice (*Oryza sativa*) and wheat (*Triticum aestivum*) given their rapid growths and thus possibility to obtain faster results (Figure 5). Garden cress (*Vigna radiata*) was also significantly studied for germination trials due to fast

germination and development of only the primary root. Microalgal/cyanobacterial derived products are a relatively unexplored resource (when compared with seaweeds), therefore there is a great opportunity in this field to develop novel research, more field trials in crop productivity, in natural environmental conditions as well as in stress abiotic conditions (e.g., salinity, temperature, and drought).

## Biofertilizers: improvement of soil quality

Some microalgal and cyanobacterial species are also well-known for their positive effect on soil quality, given their ability to fix atmospheric nitrogen, increase the availability of essential nutrients or improve soil physical and chemical properties. Furthermore, as photosynthetic microorganisms, microalgae and cyanobacteria can recover nutrients from different sources, therefore reducing the use of traditional fertilizers.

## Nitrogen fixation

As previously mentioned, most research on the agricultural applications of algae has focused on the use of cyanobacteria, given their ability to fix atmospheric nitrogen to organic nitrogen forms, that can be assimilated by higher plants (Nilsson et al., 2002; Pereira et al., 2009; Prasanna et al., 2014; Zayadan et al., 2014; Buenaventura and Barrientos, 2019; Shamim et al., 2020; Bao et al., 2021; Zhang et al., 2021) (Table 1). Altogether, the use of cyanobacteria in soils has been reported to promote nutrient recycling with increased availability of nitrogen, and to promote efficient C-N sequestration in the soil alongside the enrichment of the crop's micronutrients fraction (Swarnalakshmi et al., 2013;




FIGURE 5

Distribution of studies regarding the application of microalgae in agriculture, by the target plants.

Prasanna et al., 2015b; Prasanna et al., 2015c). When applied directly to the soil, cyanobacteria (alone or in consortia) can improve nitrogen availability and form soil crusts that avoid nitrogen leaching. This significantly improves plant growth whilst economizing fertilizers by 25–30% and minimizing the risks of water contamination (Pereira et al., 2009; Buenaventura and Barrientos, 2019; Ramírez-López et al., 2019; Bao et al., 2021; Zhang et al., 2021).

One mode of application that has been considered an economically attractive option, was the use of cyanobacteria as a matrix to develop biofilms in the soil (Prasanna et al., 2011; Prasanna et al., 2020). Prasanna et al. (2013a; 2014; 2015a; 2015b; 2015c). and Swarnalakshmi et al. (2013) have extensively used the cyanobacterium *Anabaena torulosa* to develop biofilms, due to its inherent characteristics, such as the presence of an extensive nutrient-rich mucilage and its facultative mode of nutrition, which can improve the supply of N, P and C to the plant. Moreover, the cyanobacterial biofilms can further enhance plant growth through the production of phytohormones (i.e. auxins, gibberellins, cytokinins), bioactive metabolites (i.e. aurilide, barbamide, garbamide), and increased content of glomalin-related soil protein in the soil, as observed for several crops (Prasanna et al., 2013a; Prasanna et al., 2013b; Prasanna et al., 2014; Prasanna et al., 2015b).

Another cyanobacteria that was widely used as a nitrogen-fixing biofertilizer was *Nostoc* (Table 1). Nilsson et al. (2002) screened various *Nostoc* isolates for their ability to associate with rice, which promoted higher  $N_2$  fixation rates compared to free-living cyanobacteria. In another study, *N. muscorum* and *N. rivulare*, either alone or in combination with N-fertilizer, were also shown to associate with soybean, resulting in a significant increase in plant height, leaf area, and fresh weight (Sholkamy et al., 2015). Ultimately, Buenaventura and Barrientos (2019) demonstrated that the supplementation with *N. commune* allowed for the reduction of the recommended dose of inorganic fertilizer by half since it provided a similar nitrogen content to the inorganic

fertilizer, where similar plant height and harvest index were achieved.

The use of microalgae as nitrogen-fixing biofertilizers has also been studied. Different methods of application of microalgal monocultures of *Microcystis aeruginosa* and *Chlorella* sp. were studied on maize (Grzesik and Romanowska-Duda, 2014) and willow plants (Grzesik et al., 2017). The authors concluded that applying the microalgae to grains prior to sowing, was more profitable than continuously applying them to roots *via* the substrate. In addition, Grzesik et al. (2017) studied the foliar application of the referred monocultures, from which the physiological performances and growths of plants were significantly enhanced. In both studies, the increased height, number of shoots, and length of plants, could be related to the accumulation of higher quantities of active compounds from the microalgae strains. Moreover, the applied monocultures improved the NPK content of plants, the enzyme activity (dehydrogenases, RNase, acid or alkaline phosphatase and nitrate reductase), as well as the overall photosynthesis process (stability of cytomembranes, chlorophyll content, transpiration, stomatal conductance and reduced intercellular  $CO_2$  concentration) (Grzesik and Romanowska-Duda, 2014; Grzesik et al., 2017).

Finally, the use of consortia with both cyanobacteria and microalgae has also been explored. Ramírez-López et al. (2019) showed that it was possible to reduce 75% the chemical fertilization for wheat (*Triticum aestivum* L.), due to the complementary effect of a photosynthetic and N-fixing microbial consortium, including cyanobacteria and Chlorophyta species, that had beneficial effects on plant growth and the soil. The same was evidenced by Dineshkumar et al. (2018), where reducing the chemical N fertilizer up to 50 or 75% of the recommended dose while complementing with the application of microalgae (*C. vulgaris* and *Arthrospira platensis*), showed better results than merely using the recommended N dose. In conclusion, the use of microalgae and cyanobacteria as biofertilizers has shown promising results in promoting nutrient recycling, improving



TABLE 1 Impacts of microalgae and cyanobacteria on soil quality regarding their nitrogen fixation ability.

Microalgae/Cyanobacteria	Target plant	Results	Reference
<i>Anabaena variabilis</i>	Rice	Reduction of 50% in N fertilizer with increased rice yields up to 22% compared with conventional fertilizer treatment	(Bao et al., 2021)
<i>Anabaena azotica</i>	Rice	30% replacement of N fertilizer with cyanobacteria could sustain or improve rice yield	(Zhang et al., 2021)
<i>Anabaena torulosa</i>	Wheat	Similar/higher contents of available NP with single or combined inoculation with bacteria compared to chemical controls	(Swarnalakshmi et al., 2013)
<i>Anabaena</i> spp., <i>Nostoc</i> spp. <i>Anabaena</i> -based biofilms	Rice	Better nutrient availability in the soil and uptake in plants, and increments in grain N and P	(Prasanna et al., 2015a)
	Maize	Enhanced N availability in the soil due to the application of cyanobacterial formulations	(Prasanna et al., 2015c)
	Mung bean and soybean	The <i>Anabaena</i> -based biofilm showed promising results for soybean with 12–25% enhanced yield and microbial activity	(Prasanna et al., 2014)
	Wheat	Significant enhancement in microbial activity and plant growth/yields and savings of 25% N in the wheat–rice cropping sequence.	(Prasanna et al., 2013a)
<i>Anabaena iyengarii</i> , <i>Nostoc</i> spp.	Rice	50% reduction in the use of nitrogen synthetic fertilizer (50 kg N ha <sup>-1</sup> ), resulting in the same grain yield (7.4 t ha <sup>-1</sup> ) and quality related to the fertilized control	(Pereira et al., 2009)
<i>Anabaena</i> spp., <i>Nostoc</i> spp., <i>Calothrix elenkinii</i>	Cotton	Composts fortified with cyanobacteria increased the available nitrogen in soil by 20–50%, improving germination, plant weight, and microbiological activity by 10–15%	(Prasanna et al., 2015b)
<i>Nostoc muscorum</i>	Rice	Al-acclimatized immobilized <i>N. muscorum</i> increased the growth of the rice seedlings due to augmented ammonia excretion	(Shamim et al., 2020)
<i>Nostoc commune</i>	Rice	Treated seeds with half fertilizer dosage generated similar plant height, number of tillers, and harvest index compared to full fertilizer, providing similar nitrogen content	(Buenaventura and Barrientos, 2019)
<i>Nostoc</i> spp.	Rice	Cyanobacteria association fixed more N <sub>2</sub> than when in free-living form	(Nilsson et al., 2002)
	Soybean	Improvement of plant growth of plants treated with <i>Nostoc</i> spp. alone or combined with N-fertilizer, allowing a 50% reduction in fertilizers	(Sholkamy et al., 2015)
<i>Nostoc</i> sp., <i>Calothrix ghosei</i> , <i>Hapalosiphon intricatus</i>	Wheat	Cyanobacterial isolates along with a 1/3 N fertilizer dose allowed a comparable grain yield to a full fertilizer dose	(Karthikeyan et al., 2007)
<i>Nostoc entophytum</i> , <i>Oscillatoria angustissima</i>	Pea	Biofertilization combined with half fertilizer dose was more effective than the addition of the full dose	(Osman et al., 2010)
<i>Anabaena variabilis</i> , <i>Chlorella vulgaris</i>	Rice	10–20% increase in germination rate and plant length	(Zayadan et al., 2014)
<i>Anabaena</i> sp., <i>Chlorella</i> sp., <i>Microcystis aeruginosa</i>	Maize	Increase of germination and growth of maize seedlings and intensification of metabolic processes	(Grzesik and Romanowska-Duda, 2014)
	Willow	Enhancement of plant growth and health and improvement of chlorophyll content and gas exchanges in leaves	(Grzesik et al., 2017)
<i>Aphanizomenon aphanizomenoides</i> , <i>Leptolyngbya</i> sp., <i>Anabaena oscillarioides</i> , <i>Monoraphidium</i> sp., <i>Chlorella</i> sp.	Wheat	Reduction of 75% of chemical fertilization for wheat production	(Ramírez-López et al., 2019)
<i>Arthrospira platensis</i> , <i>Chlorella vulgaris</i>	Rice	Higher nutrient availability in soil and plant leaves	(Dineshkumar et al., 2018)
<i>Phormidium</i> sp.	Wheat	<i>Phormidium</i> sp. was shown to solubilize phosphate, fix atmospheric nitrogen, and produce hydrogen cyanide.	(Mazhar and Hasnain, 2011)

nitrogen availability, and enhancing plant growth while reducing the need for inorganic fertilizers, therefore holding great potential for sustainable agriculture.

## Nutrient availability

In the past three decades there have been secondary micronutrient deficiencies in the soils, due to the imbalanced use of macronutrient fertilizers, decreased use of organic manure, reduced recycling of crop residues, and bumper harvests. Microalgae and cyanobacteria are photosynthetic microorganisms that can uptake and store essential nutrients (N and P) in their biomass, even from sites where they are scarce. In addition, they can accumulate other important micronutrients like potassium, magnesium, sulfur, and iron, which are involved in redox

reactions in the plant metabolism (Ronga et al., 2019; Gonçalves, 2021). Thus, as it has been reported in the literature (Table 2), microalgae or cyanobacteria can be considered as an alternative source of essential macro- and micronutrients for plants' growth.

In addition to their nitrogen-fixation capacity, cyanobacteria-bacteria biofilms or consortia can improve nutrient availability in the soils (Rana et al., 2012; Manjunath et al., 2016; Simranjit et al., 2019). Simranjit et al. (2019) showed that the application of these biofilms improved the composition of the soil, not only in N (50–90%) but also in P and organic C by 40–60%. Moreover, these biofilms improved the bioavailability of micronutrients such as Cu, Fe, Mn, and Zn, which reflected a positive effect on the leaves photosynthetic pigments. Manjunath et al. (2016) also observed improvements in the bioavailability of soil macro/micronutrients, by applying *Anabaena* in microbial biofilms or consortia, and *Calothrix* sp. The synergistic action of bacteria and cyanobacteria

TABLE 2 Impacts of microalgae and cyanobacteria on soil quality regarding nutrient availability.

Microalgae/ Cyanobacteria	Target plant	Results	Reference
<i>Anabaena</i> -based biofilms	Cucumber	Increase of N by 50–90%, P, and organic C by 40–60% in soil, over control Significant augmentation in leaf chlorophyll, soil microbiological parameters, and nutrient bio-availabilities (Zn, Mn, Cu, and Fe)	(Simranjit et al., 2019)
<i>Anabaena</i> -based biofilms <i>Calothrix</i> sp.	Okra	Increase in Zn and Fe contents of soil, leading to improved root yield and weight	(Manjunath et al., 2016)
<i>Anabaena</i> spp., <i>Calothrix</i> sp.	Wheat	Biofortification of wheat plants with micronutrients (Fe, Zn, Mn, and Cu), leading to 50% savings of N fertilizer without compromising grain quality	(Rana et al., 2012)
<i>Arthrospira platensis</i>	Strawberry	Increased micronutrients (Fe and Si) in the root system	(Soppelsa et al., 2019)
<i>Arthrospira platensis</i> , <i>Chlorella vulgaris</i>	Mung bean, onion, maize	Higher nutrient availability in soil and plant leaves and increase in soil microflora and weak carbon dioxide emission in the treatments	(Dineshkumar et al., 2019; Dineshkumar et al., 2020a; Dineshkumar et al., 2020b)
<i>Chlorella</i> sp., Bioguan (guano, macroalgae, and <i>Arthrospira platensis</i> )	Barley	Higher nutrient uptake efficiencies using bioguan	(Atzori et al., 2020)
<i>Chlorella vulgaris</i>	Black gram	Higher nutrient availability in soil	(Dineshkumar et al., 2020c)
	Wheat	Similar growth of microalga-treated plants with conventional fertilizer application	(Schreiber et al., 2018)
	Lettuce	Improvement of plant nutrients, which enhanced all the physiological reactions that led to a good growth	(Faheed and Fattah, 2008)
	Maize	Improvement of soil physical and chemical characteristics, plant nutrient status, saving the addition of secondary and microelements	(Shaaban, 2001b)
<i>Chlorella sorokiniana</i>	Wheat	Improvement of soil fertility and plant nutrition	(Kholssi et al., 2019)
<i>Chlorella variabilis</i> , <i>Scenedesmus obliquus</i>	Maize, soybean	Improvement of nitrogen and phosphorus content	(Loganathan et al., 2020)
<i>Chlorella</i> sp., <i>Neochloris conjuncta</i> , <i>Botryococcus braunii</i>	Maize	Slight increase in plant weight and nutrient uptake using digestates containing <i>Chlorella</i> sp. at the lowest dose	(Ekinci et al., 2019)
<i>Acutodesmus dimorphus</i>	Tomato	Increase in plant growth	(Garcia-Gonzalez and Sommerfeld, 2016)
<i>Chlorococcum</i> sp.	Mung bean, tomato, cucumber, pepper	20% biomass showed a better response in growth parameters such as root length, shoot length, number of lateral roots, and number of leaves	(Deepika and MubarakAli, 2020)
<i>Tetraselmis</i> sp.	Date palm	Increase of soil NPK content	(Saadaoui et al., 2019)

increased Fe and Zn contents, contributing to an improved root yield and weight of okra plants. These two micronutrients are amongst the most limiting nutrients for plant growth. However, since several cyanobacteria can sequester Fe or Zn from metallothioneins, they can be used for biofortification to cover these crop deficiencies (Rana et al., 2012; Manjunath et al., 2016).

Regarding the application of microalgal species, *Chlorella vulgaris* has been a widely studied species on different plants (Shaaban, 2001b; Faheed and Fattah, 2008; Dineshkumar et al., 2020c). In a study involving the use of both cyanobacteria and microalgae, Dineshkumar et al. (2019; 2020b) tested *C. vulgaris* and *A. platensis* mixed with cow manure as a biofertilizer for maize and onion. For maize, all the treatments recorded significantly improved the levels of N, Mn, P, and K. In contrast, low levels were observed for Ca, Zn, Fe, and CO<sub>3</sub> and Na contents. The combination of manure with microalgae allowed for the best growth of maize during the early stages, specifically up to 51.1% after 60 days of planting (Dineshkumar et al., 2019). The same trend was observed when applying identical conditions to onion plants. The microalgal treatments mixed with cow manure allowed for the maximum micro and macronutrient availability, which yielded the best growth parameters in onion plants. All treatments originated onions larger than the marketable size, with improved biochemical (e.g., soluble sugars, phenols, free amino acids, and indoles) and mineral compositions (e.g., Na, K, P, Ca, Fe, Mg, and Mn), that result in increased market value. The same group also tested the foliar application of both species in mung bean plants (Dineshkumar et al., 2020a). Likewise, plants treated with *C. vulgaris* and *A. platensis* generated superior levels on the composition of mung bean leaves and soil and improved the physical characteristics of green gram including water absorption index, water solubility index, and water and oil absorption. Recently, Dineshkumar et al. (2020c) concluded that *C. vulgaris* extracts could be used as an eco-friendly and potentially economically viable foliar spray treatment for black gram (*Vigna mungo* L.). The soil properties and production yields were improved, as well as fortified the seed flour with beneficial minerals for consumers, such as Na, K, Ca, Mg, and P.

Nutrient depletion is a major problem for crop production, given the insufficiency of direct sunlight and continuous cultivation with intensive use of fertilizers and pesticides. As previously stated, microalgae can recover nutrients from sites where nutrient availability is limited. Schreiber et al. (2018) compared the application of mineral and *Chlorella vulgaris* fertilizers in nutrient-poor soil substrates. For all experimental conditions, the biofertilization enhanced the growth of wheat plants in comparison to the unfertilized controls (nutrient limitation). No significant differences were observed between mineral and *Chlorella* fertilizers. When compared with wheat plants grown without nutrient limitation, the fertilization with *Chlorella vulgaris* facilitated the growth of wheat with similar nitrogen contents, but lower phosphorus levels in plant tissues. This suggests that microalgae cells can cover the nutrient deficiency of the soils, though, while N can be released in a form that wheat roots can easily uptake, P might not, or is released slowly. Nonetheless, the root-hair properties of plants were similar among treatments, which confirms the

conversion of microalgal P to plant-available forms in the soil. These results validate that microalgal biomass can support crop growth on marginal soils, with similar performance to mineral fertilizers.

The use of other microalgal species has also been evaluated, as is the case of *Acutodesmus dimorphus*, whose culture, extracts, and dry biomass were tested in seed priming, foliar spraying, and biofertilization of tomato plants (Garcia-Gonzalez and Sommerfeld, 2016). The authors were able to achieve early germination with the seed treatment, as well as longer plants, a higher number of flowers, and braces per plant, with foliar application of the extract. Furthermore, the application of dry *A. dimorphus* biomass 22 days prior to transplant, significantly enhanced plant growth, when compared to the application at the time of transplant. These results suggest that a preliminary application is required for the biomass to break down and increase the nutrient availability for plant uptake. In another case, Saadaoui et al. (2019) studied the effects of fertilizing date palms with *Tetraselmis* sp. The authors observed that after 3 months of treatment, the supplementation with 0.5 g of *Tetraselmis* sp. increased the NPK content of soil and maintained a safe level of heavy metals. This led to higher plant growth rates of date palm when compared to conventional fertilizer, with 100% survival rates, greater number of leaves ( $3.17 \pm 0.14$  vs.  $3 \pm 0.66$ ), high ability to root largest stem thickness, longer shoot, and higher chlorophyll content.

## Soil amendment

Microalgae and cyanobacteria can also enhance soil properties (aggregation, porosity, permeability, ventilation, and humidity) (Sharma et al., 2012; Baweja et al., 2019; Pan et al., 2019). As previously stated, some microalgae and cyanobacteria species can excrete polysaccharides that form an adhesive and gelatinous mesh, which binds soil particles on their surface, therefore improving water retention and soil aeration (Xu et al., 2013). This ability plays a significant role in pH and temperature regulation, as well as protection against erosion (Sharma et al., 2012; Baweja et al., 2019). These improvements positively affect crop productivity since they stimulate root growth and soil microbial activity. Therefore, microalgae and cyanobacteria can help build soil fertility (Nisha et al., 2007; Saadatnia and Riahi, 2009; Prasanna et al., 2016a; Marks et al., 2017; Barone et al., 2019b; Lv et al., 2020), prevent erosion (Hu et al., 2002; Issa et al., 2007; Kheirfam et al., 2017), recover damaged soil crusts (Acea et al., 2001), or sequester toxic pollutants from soil (Tripathi et al., 2008; Priya et al., 2014; Decesaro et al., 2016) (Table 3). Furthermore, microalgae and cyanobacteria can form associations with plant roots which are mutually beneficial. They benefit plants by producing extracellular enzymes that break down organic matter in the soil, and releasing nutrients that are essential for plant growth, such as nitrogen and phosphorus. They can also produce growth-promoting hormones that stimulate plant growth (Prasanna et al., 2011). In turn, plant roots release exudates (e.g., simple sugars, amino acids, organic acids, and other compounds) that provide a food source for the microorganisms.

TABLE 3 Impacts of microalgae and cyanobacteria on soil bioremediation.

Microalgae/Cyanobacteria	Target plant/soil	Results	Reference
<i>Chlorella</i> sp.	Agricultural soil	Increase of eukaryotic and prokaryotic biomass and the activities of heterotrophic microorganisms in the soil	(Marks et al., 2017)
<i>Chlorella vulgaris</i>	Rice in As contaminated soil	Reduction of As availability	(Srivastava et al., 2018)
	Maize and wheat	Improvement of the amount of soil organic matter and water retention capacity	(Uysal et al., 2015)
<i>Calothrix elenkinii</i>	Rice rhizosphere	Significant increases in plant growth attributes, nitrogenase activity, indole acetic acid production, and activities of hydrolytic and defense enzymes	(Priya et al., 2015)
<i>Scenedesmus quadricauda</i> , <i>Anabaena circinalis</i>	Cucumber rhizosphere	Enhancement of the rhizosphere microbial diversity	(Lv et al., 2020)
<i>Chlorella vulgaris</i> , <i>Scenedesmus quadricauda</i>	Soil cultivated with tomato	Improvement of soil biological activity	(Barone et al., 2019b)
<i>Oscillatoria</i> sp., <i>Nostoc</i> spp., <i>Scytonema</i> sp.	Burned soils	Improvement of crust formation and restoration of microbial populations	(Acea et al., 2001)
<i>Nostoc</i> sp., <i>Oscillatoria</i> sp., <i>Lyngbya</i> sp.	Erosion-prone soils	Increase of the total organic nitrogen and total organic carbon and soil organic matter	(Kheirfam et al., 2017)
<i>Chlorella sorokiniana</i>	Calcareous soils	Autotrophic microorganism development was 3.5 times higher in treatments with algae application as measured by chlorophyll pigment concentration	(Marks et al., 2019)
<i>Anabaena doliolum</i> , <i>Cylindrospermum sphaerica</i> , <i>Nostoc calcicola</i>	Pearl millet and wheat	Decline in bulk density and increase in water holding capacity, hydraulic conductivity, and mean weight diameter	(Nisha et al., 2007)
<i>Anabaena</i> spp., <i>Nostoc</i> spp.	Maize rhizosphere	Improvement of soil functional activities	(Prasanna et al., 2016a)
<i>Anabaena</i> spp.	Rice	Earlier germination and longer seedlings Enhancement of soil moisture (20%), and porosity (28%); decrease of soil bulk (9.8%) and particle density (4.8%) Increase in plant height (53%), root length (66%), fresh (69%), and dry weight (137.5%)	(Saadatnia and Riahi, 2009)
<i>Nostoc</i> sp., <i>Anabaena doliolum</i> , <i>Calothrix</i> sp., <i>Westiellopsis</i> sp., <i>Phormidium papyraceum</i>	Rice	Improvement of plant growth, yield, and mineral composition while reducing the nitrogen fertilizers	(Tripathi et al., 2008)
<i>Arthrospira platensis</i>	Diesel and contaminated biodiesel-contaminated soils	The phycocyanin extracts were more effective in biodiesel removal (88.8%) while <i>A. platensis</i> cells were better in diesel removal (63.9%)	(Decesaro et al., 2016)

These microorganisms can then form colonies around the plant roots, creating a biofilm that can help protect the roots from pathogens and other stressors, while also improving the soil structure. The improved soil structure allows for better water and nutrient retention, which further supports plant growth (Nisha et al., 2018).

In a study, a multi-strain biofertilizer (dried paste) consisting of three indigenous cyanobacterial isolates (*Anabaena doliolum*, *Cylindrospermum sphaerica*, and *Nostoc calcicole*), was applied to an organically poor semi-arid soil under limited-water conditions (Nisha et al., 2007). The native strains improved carbon and nitrogen mineralization, by promoting soil microbial activities and decreasing the C:N ratio. Also, a decline in bulk density and an increase in water holding capacity and hydraulic conductivity of soil were observed. Plant growth and yield of pearl millet and wheat increased in response to cyanobacterial biofertilizer, which was more pronounced at lower water levels (6%). In another study, microalgal suspensions of alive *S. quadricauda* and *Anabaena*

*circinalis* were evaluated in cucumber cultivation. The application of a high concentration of *S. quadricauda*, in the soil, boosted the diversity of rhizosphere fungi of cucumber, especially of growth-promoting bacteria and fungi (*Azotobacter*, *Bacillus*, *Pseudomonas*, *Cryptococcus*, *Fusarium*, *Penicillium*, and *Trichoderma*), which resulted in increased height, number of leaves, flower buds, and stem diameter of cucumber (Lv et al., 2020).

Enriching the biological properties of soil crusts, by increasing its microbial population, can be an effective strategy to improve soil chemical properties. Moreover, because of their filamentous structure and their ability to secrete polysaccharides and other cementing substances (amongst others), cyanobacteria can generate a prosperous environment for microbial populations to proliferate and form a crust, which has a significant role in preventing soil erosion, improving water kinetics and plant growth (Nisha et al., 2018).

Accordingly, Acea et al. (2001) investigated the potential value of cyanobacterial suspensions (*Nostoc*, *Oscillatoria* and *Scytonema* species) for accelerating soil recolonization and promoting

microbiotic crust formation after a fire. The heated soils inoculated with cyanobacteria were quickly colonized and, after 2 months, the cyanobacterial filaments, and associated fungal hyphae, built up a matrix that aggregated surface soil particles, therefore improving crust formation. These crusts were not observed in the uninoculated soils. In another study, Kheirfam et al. (2017) showed that the inoculation of cyanobacteria suspensions (*Nostoc*, *Oscillatoria*, and *Lyngbya* species) had a more effective role in N fixing (240%) and organic matter storing (40%), compared to the other inoculation treatments, which ultimately improves soil quality to prevent soil degradation.

Finally, microalgae and cyanobacteria can perform the remediation of soils by removing heavy metals (e.g., cadmium, lead, and chromium), trace elements (e.g., iron, zinc, copper, and manganese), hydrocarbons, and others, which helps build soil quality and fertility (Priya et al., 2015; Suresh Kumar et al., 2015). Heavy metals can be transformed, detoxified, and volatilized by microalgae, and since microalgae are non-pathogenic there is no risk of accidental release into the atmosphere (Priya et al., 2015). Microalgae degrade the pollutants enzymatically but can also absorb them onto their surfaces due to the high metal binding capacity of polysaccharides, proteins, or lipids present on their cell walls (Suresh Kumar et al., 2015). This capacity is crucial in rice production, since an increase in land contamination has been reported, which not only reduces arable land but could negatively impact human health due to heavy metal accumulation in rice grains (Tripathi et al., 2008; Srivastava et al., 2018). The inoculation of cyanobacteria slightly declined Cd, Ni and As in the content of rice plants, while reducing the nitrogen requirements by 30 kg/ha. Aside from enhancing growth, it also fortified the composition of essential micronutrients while maintaining the level of toxic metals under safe limits (Tripathi et al., 2008). A combination of *C. vulgaris* and *Pseudomonas putida* mitigated the As stress during P-enriched conditions by reducing As availability, as well and modulating its uptake and detoxification mechanisms in rice plants (Srivastava et al., 2018).

## Plant growth stimulation

The use of microalgae and cyanobacteria (and/or their extracts) can directly stimulate plants' growth and development by improving germination rates and plant characteristics, such as shoot and root length, leaf area, and nutrient contents. These enhancements are accomplished due to microalgal/cyanobacterial metabolites (e.g., phytohormones, amino acids, vitamins, polysaccharides, polyamines, etc.), which can trigger several metabolic responses, such as respiration, photosynthesis, nucleic acid synthesis, chlorophyll production and ions uptake (Chiaiese et al., 2018; Chanda et al., 2019). Table 4 demonstrates examples of microalgae and cyanobacteria (biomass and/or extracts) that have been studied for their direct stimulation of higher plants growth.

The potential of *C. sorokiniana* to be used as substitute for chemical fertilizers, and its capacity to stimulate wheat plants was studied by Kholssi et al. (2019). Apart from the harvested biomass, resuspended in fresh and spent medium, the authors also analyzed the filtered medium where the microalga was cultivated and compared it

with fresh medium alone (control). All treatments resulted in a higher germination percentage than the control, however, only the treatments containing spent medium (with or without microalga) were statistically different from the control. A general enhancement of plant growth was observed with the microalga treatments, although the most significant increase in plant length (30%) and fresh weight (77%) was obtained with the filtrate of *C. sorokiniana* (spent medium), which suggests that the microalga excreted beneficial compounds for plant growth into the medium. From a biorefinery perspective, these results are extremely interesting, since the supernatant has the potential to be used for the biostimulation of plants, whilst the harvested biomass can be used for other high-value applications like food and/or feed (Ferreira et al., 2022).

The biostimulant potential of *Chlorella vulgaris* (auto and heterotrophic) was addressed by Uysal et al. (2015), where an increase in height was observed for wheat and barley plants when compared with the control. Barone et al. (2019a) reported a positive effect on the biomass productivity of tomato plants and microalgae in a co-cultivation system, especially using *Scenedesmus quadricauda* biomass in the presence of digestates from the waste of an agrolivestock farm. The highest weight values of fresh plant shoot and dry whole plant were observed in the co-cultivation system with only *C. vulgaris*, where a 2.5-fold increase was obtained over the control. Barone et al. (2018) also found that *C. vulgaris* and *S. quadricauda* extracts had biostimulant effects on the expression of root traits and genes related to the nutrient acquisition in sugar beet (genes putatively involved in sulfate starvation). In a similar study, Puglisi et al. (2020) tested *S. quadricauda* extracts, demonstrating a beneficial impact on the growth of lettuce seedlings, especially at the shoot level, along with increased contents of dry matter, chlorophylls, carotenoids, and protein. Moreover, these extracts were also proven to positively influence the activities of various enzymes involved in the carbon and nitrogen primary metabolisms.

Plaza et al. (2018) evaluated the effects of foliar application of *Scenedesmus* sp. and *A. platensis* hydrolysates in petunia plants. The application of *Scenedesmus* improved the plant nutrient status, and accelerated plant development and the flowering process, while *Arthrospira* improved the root dry matter, the number of flowers, and the water content.

The application of various dosages of *Arthrospira* in radish plants was also explored by Godlewska et al. (2019). An increase in length and fresh weight was observed for homogenate treatments of seeds and of filtrate as a foliar spray. The highest increase in chlorophyll was observed for lower concentrations. It is also relevant to acknowledge that the lowest dosages of *Arthrospira* always achieved better results compared with the commercial biostimulant. Akgül (2019) demonstrated that up to 75% microalgal cell extract promoted an increase in germination and seedling of wheat and barley plants and that higher concentrations had an inhibitory effect. Michalak et al. (2016) performed field trials to evaluate the effect of supercritical extracts of *A. platensis* in winter wheat plants. The best results were achieved by *A. platensis* extracts (higher number of grains in ear and shank length). Moreover, these extracts showed similar biostimulant properties to commercial products, with the additional advantage of being a natural and environmentally friendly source of biologically active compounds.



TABLE 4 Impacts of microalgae and cyanobacteria on plant growth stimulation (biostimulant).

Microalgae	Target plant	Results	Reference
<i>Chlorella sorokiniana</i>	Wheat	Significant increase in plant length and dry weight of wheat plants using the spent medium after culture separation	(Kholssi et al., 2019)
	Rice and tomato	The presence of phytohormones in algal extract triggered earlier seed germination at higher than 40% (v/v)	(Do et al., 2020)
	Orchid	Explants supplemented with microalga suspension, or the supernatant showed similar growth to the control	(Pereira et al., 2018)
<i>Chlorella vulgaris</i>	Cress, mung bean, and wheat	Increase of germination index of cress seeds using biomass extracts of <i>C. vulgaris</i> grown in synthetic medium	(Ranglová et al., 2021)
	Mung bean, rocket, and cress	Increase in radicle and plumule length with aqueous extract treatment	(Al dayel et al., 2020)
	Wheat and French bean	Polysaccharides solutions (3 and 5 mg mL <sup>-1</sup> ) extracted from mixotrophic cultures enhanced seedling growth (root length, leaf area, shoot length, photosynthetic pigments, protein, and carbohydrates), due to increased antioxidant activity	(El-Naggar et al., 2005)
	Wheat	Alga extract at 50% (v/v) increased fresh weight (60.7%), and yield (>140%), while 100% (v/v) increased dry weight (95%), spikes weight (60%), and grain weight (160%).	(Shaaban, 2001a)
<i>Chlorella vulgaris</i> , <i>Scenedesmus quadricauda</i>	Tomato	Increase in productivity and plant growth (fresh and dry weight), in a co-cultivation system with tomato and microalgae	(Barone et al., 2019a; Barone et al., 2019b)
	Sugar beet	Increase in expression of root traits and genes related to nutrient acquisition with microalgae extracts	(Barone et al., 2018)
<i>Scenedesmus quadricauda</i>	Lettuce	Treatment with microalgae extract stimulated the primary metabolisms, improving the plant growth and contents of dry matter, pigments, and protein	(Puglisi et al., 2020)
<i>Scenedesmus</i> sp., <i>Arthrospira platensis</i>	Petunia	Foliar spraying with microalgae hydrolysates generated higher rates of root growth, leaf and shoot development, and early flowering	(Plaza et al., 2018)
<i>Chlorella vulgaris</i> , <i>Arthrospira platensis</i>	Mung bean	<i>Arthrospira</i> extracts increased protein, NPK, Ca, Mg, and Zn contents, and <i>Chlorella</i> extracts boosted the amino acids	(Dineshkumar et al., 2020a)
<i>Arthrospira platensis</i>	Radish	Microalgae filtrates and homogenates increased plant length, fresh weight, and chlorophyll content	(Godlewska et al., 2019)
	Wheat and barley	Promotion of germination and seedling growth at concentrations up to 75% microalgal extract	(Akgül, 2019)
	Wheat	A higher number of grains in ear and shank length, at 1.5 L/ha of supercritical microalgal fluid extraction	(Michalak et al., 2016)
	Strawberry	Microalgal hydrolysate improved biomass accumulation (iron and silicate content) in roots	(Soppelsa et al., 2019)
	Lettuce	Microalgal foliar application improved the quality characteristics (soluble solids, titratable acidity, ascorbic acid, chlorophyll) of post-harvested and stored lettuce	(Silva et al., 2017)
	Cucumber, mung bean, lettuce, and beet	Foliar application of microalgae hydrolysates increased the growth of seedlings and spermidine content of lettuce and increased the fresh weight of cucumber cotyledons. Foliar application of aqueous suspension of biomass on beet seedlings improved growth due to enhanced chlorophyll synthesis and amino acid and protein increments	(Mógor et al., 2018a; Mógor et al., 2018b)
	Eggplant	Foliar application (10 g/L) increased fruit yield and pulp firmness after 6 days of storing, while higher concentrations stimulated vegetative growth and reduced plant yield	(Dias et al., 2016)
	Pepper and tomato	Spraying with total polysaccharides extract increased plant size, roots weight, and size and number of nodes	(El Arroussi et al., 2016)
	Wheat	Supercritical fluid extracts were applied obtaining similar plant height, ear length, and shank length to the control	(Michalak et al., 2016)
<i>Nannochloropsis oculata</i>	Tomato	Pulverized microalgal biomass promoted similar growth to the control, but with improved fruit quality (sugar and carotenoids)	(Coppens et al., 2016)

(Continued)

TABLE 4 Continued

Microalgae	Target plant	Results	Reference
<i>Acutodesmus dimorphus</i>	Tomato	Treatments with microalgal culture, culture medium, and aqueous extracts improved seed germination. Foliar pulverization of alga extracts increased plant height and number of flowers and branches per plant	(Garcia-Gonzalez and Sommerfeld, 2016)
<i>Desmodesmus subspicatus</i>	Orchid	Microalgal aqueous cell extract increased germination and shoot formation	(Navarro et al., 2021)
	Mung bean and onion	Aqueous microalga suspensions improved bulb caliber and yield and incremented sugars and protein in bulbs	(Gemin et al., 2019)
<i>Messastrum gracile</i> , <i>Chlorella vulgaris</i>	Orchid	Microalgal aqueous extracts improved elongation, root formation, and plant survival	(Corbellini et al., 2020)
<i>Chlorella vulgaris</i> , <i>Chlorella sorokiniana</i> , <i>Chlamydomonas reinhardtii</i> , <i>Dunaliella salina</i>	Tomato	Injection of microalgal polysaccharides improved the plant defense mechanisms and increased $\beta$ -1,3-glucanase activity and PUFAs contents	(Farid et al., 2019)
<i>Anabaena variabilis</i> , <i>Nostoc calcicole</i>	Corn, sorghum, rice, cowpea, kodo millet	Microalgal extracts improved germination and root, shoot, and total seedling lengths	(Suresh et al., 2019)
<i>Dunaliella salina</i> , <i>Phaeodactylum tricornutum</i> , <i>Porphyridium</i> sp., <i>Desmodesmus</i> sp., <i>Arthrospira platensis</i>	Tomato	Tomato plant leaves treated with extracted microalgal polysaccharides showed improved enzymatic activity	(Rachidi et al., 2021)
<i>Calothrix elenkini</i> , <i>Anabaena laxa</i> , <i>Anabaena torulosa</i>	Radish, wheat, and corn	Cyanobacterial extracts resulted in marginal increases in seed germination of maize, radish, and wheat	(Prasanna et al., 2020)
<i>Chlorella</i> sp., <i>Chlorella vulgaris</i> , <i>Scenedesmus acutus</i>	Mung bean	Aqueous extracts increased plant rooting with <i>Chlorella</i> treatments	(Stirk et al., 2020)
<i>Anabaena</i> sp., <i>Calothrix</i> sp., <i>Phormidium</i> sp., <i>Oscillatoria</i> sp.	Tobacco	Extracellular extract of culture filtrate promoted germination and root multiplication	(Boopathi et al., 2013)
<i>Anabaena</i> sp., <i>Oscillatoria</i> sp., <i>Phormidium</i> sp., <i>Chroococcidiopsis</i> sp., <i>Synechocystis</i> sp.	Wheat	Seed soaking in cyanobacterial suspensions increased germination, shoot length, tillering, number of lateral roots, spike length, and grain weight	(Hussain and Hasnain, 2011)
<i>Anabaena vaginicola</i> , <i>Nostoc calcicole</i>	Squash, cucumber, and tomato	Soil spraying with algal extracts increased plant height, root length, dry and fresh weight, and leaf number	(Shariatmadari et al., 2013)
<i>Phormidium tenue</i>	Korshinsk pea shrub	Algal polysaccharides extract increased germination, nutrient and carbohydrate contents, photosynthetic activity, and antioxidant activity	(Xu et al., 2013)
<i>Scytonema bohnerei</i> , <i>Calothrix</i> sp., <i>Nostoc</i> sp., <i>Anabaena</i> sp., <i>Dolichospermum spiroides</i> , <i>Aphanothece stagnina</i>	Pea and cucumber	The cyanobacterial extract increased radicle, plumule, and total seedling length	(Gayathri et al., 2017)
Various species	Cucumber, soybean, and mung bean	Cyanobacterial extracts had cytokinin-like activity in soybean and auxin-like activity in root formation of cucumber cotyledon	(Stirk et al., 2002)
Various species	Tomato	Microalgal extracts increased root and shoot lengths, dry weight, and nutrient uptake and promoted the accumulation of palmitic acid, stearic acid, pyridine-3-carboxamide (an amide active form of vitamin B3), and linolenic acid	(Mutale-joan et al., 2020)

In conclusion, the use of microalgae and cyanobacteria as biostimulants for higher plants has shown promising results in enhancing growth and development, with their metabolites triggering several metabolic responses. These findings suggest a potential for these organisms to replace chemical fertilizers and serve as a sustainable and environmentally friendly source of biologically active compounds.

## Expansion of plant tolerance to abiotic stress

Under non-stress conditions, plants use the most energy in processes necessary for maintenance, and vegetative, and generative growth. However, when exposed to extreme environmental conditions (e.g., cold, heat, drought, salinity), plant resources are

allocated towards stress mitigation, which decreases plant growth and final yield (Table 5).

## Temperature

Extremely low or high temperatures negatively affect the metabolic activity of plants, and damage cell membranes, therefore affecting the nutritional composition of plants and the rates of photosynthesis and transpiration. To improve the plant's response to these harmful environmental conditions, a very promising option can be the use of microalgal biostimulants, as they are a rich source of antioxidants and other bioactive compounds that can modulate plant response mechanisms to cope with adverse conditions. The use of algal-based products as biostimulants has already been successfully applied in crops under temperature stress conditions (Figure 4D; Table 5).

Heat-stressed *Arabidopsis thaliana* treated with *Nostoc muscorum* exhibited significantly lower root hair death when compared to untreated seedlings (Chua et al., 2020). Moreover, they identified proline as a compound of interest responsible for stress response. Kopta et al. (2018) showed that the application of a

bacterial-algal biostimulant enhanced the fresh weight of two lettuce varieties in the spring and summer seasons. The summer crop of romaine lettuce treated with the biostimulant presented enhanced antioxidant capacity and carotenoid contents. Nonetheless, it is essential to highlight the vital role that the biostimulant application had in mitigating the temperature stress, which positively impacted the harvesting yield during the summer season.

## Salinity

An *Arthrospira platensis* hydrolysate was tested for its capacity to mitigate the negative effect of salt on *Petunia x hybrida* (Plaza et al., 2018; Bayona-Morcillo et al., 2020) and *Pelargonium hortorum* (Tejada-Ruiz et al., 2020). Both studies observed an increase in plant growth and flower development, under high salinity. *Dunaliella salina* extracts were also shown to alleviate the salt stress in tomatoes (El Arroussi et al., 2018) and pepper plants (Guzmán-Murillo et al., 2013). According to GC-MS metabolomics analysis, the exopolysaccharides of *D. salina* triggered the activation or inhibition of the metabolic pathways involved in the plant's

TABLE 5 Impacts of microalgae and cyanobacteria on the enhancement of tolerance to abiotic stress (heat, temperature, drought).

Abiotic stress	Microalgae	Target plant	Results	Reference
Heat	<i>Nostoc muscorum</i>	Thale cress	Decrease in programmed cell death in root hairs compared to untreated seedlings	(Chua et al., 2020)
	<i>Chlorella vulgaris</i>	Lettuce	Increase in fresh weight of summer lettuce plants	(Kopta et al., 2018)
Salinity	<i>Arthrospira platensis</i>	Petunia and geranium	Stimulation of vegetative growth and plant yield and promotion of flowering	(Plaza et al., 2018; Bayona-Morcillo et al., 2020; Tejada-Ruiz et al., 2020)
	<i>Nostoc calcicole</i> , <i>Anabaena variabilis</i> , <i>Nostoc linkia</i>	Rice	Increase of growth and decrease of soil electrical conductivity	(El-Sheekh et al., 2018)
	<i>Nostoc muscorum</i>	Rice	Protection of rice plants towards salt and heavy metals (CdCl <sub>2</sub> )	(Shamim et al., 2020)
	<i>Dunaliella salina</i>	Tomato	Attenuation of salt stress on plant growth by activation/inhibition of metabolic pathways involved in tolerance to stress	(El Arroussi et al., 2018)
	<i>Phaeodactylum tricornutum</i> , <i>Dunaliella salina</i>	Pepper	Mitigation of oxidative stress and increase in antioxidant enzymes	(Guzmán-Murillo et al., 2013)
	Cyanobacteria	Soybean	Improved levels of physio-biochemical attributes and antioxidant (non-enzymatic and enzymatic) defense systems	(Zaki et al., 2019)
	<i>Cyanothece</i> sp.	Barley	Improvement of germination, shoot, and root lengths, and fresh weight with the combined use of cyanobacterium, rhizobacterium, and methyl salicylate	(El Semary et al., 2020)
Drought	<i>Euglena gracilis</i>	Tomato	Earlier fruit ripening process and improvement of fruit quality (antioxidants and carbohydrates)	(Barsanti et al., 2019)
	<i>Chlorella vulgaris</i>	Guar	Increase in antioxidant activity and reduction of oxidative stress, improving shoot length, fresh/dry weight, leaf number, and area	(Kusvuran and Kusvuran, 2019)
	<i>Nostoc ellipsosporum</i> , <i>Nostoc punctiforme</i>	Pearl millet and wheat	Improved productivity under semi-arid conditions	(Nisha et al., 2018)
	<i>Arthrospira platensis</i>	Grape berry	Improvement of berry composition under drought conditions	(Salvi et al., 2020)

response to stress, such as jasmonic acid-dependent pathways. Root growth of pepper plants was favorably influenced by the microalgae treatments (*D. salina* and *Phaeodactylum tricornutum*), which increased the content of antioxidant enzymes and reduced the oxidative stress imposed by the increasing salt concentrations (Guzmán-Murillo et al., 2013). Cyanobacteria, particularly *Nostoc*, were also shown to mitigate the saline stress by decreasing the soil's electrical conductivity (El-Sheekh et al., 2018). Other studies showed that the integrative application of cyanobacteria with other microbes or antioxidant substances could also significantly improve growth characteristics under saline conditions (Zaki et al., 2019; El Semary et al., 2020).

## Drought

Drought stress has a negative impact on several cultures. Tomatoes are particularly very sensitive to this type of stress, which strongly affects photosynthesis and, consequently, plant growth and yield. To mitigate drought stress, Barsanti et al. (2019) explored the effects of  $\beta$ -(1,3)-glucan (paramylon) from the microalga *Euglena gracilis*. Paramylon-treated plants showed no signs of wilting compared to untreated plants, and while the density and length of the root system were drastically reduced, there was an increase in lateral rootlets. The water stress imposed on tomato plants negatively affected the main ecophysiological parameters of leaves (water potential, CO<sub>2</sub> assimilation, internal concentration, and stomatal conductance). However, the paramylon treatment allowed the tomato plants to recover to the values of control plants (non-stressed), after 1-2 weeks. The fruit size of paramylon-treated plants was like the ones under the optimal water regime, although the former reached the first ripening stage two weeks earlier than the latter. Moreover, the content of antioxidant compounds (carotenoids, phenolic acid, and vitamins), and soluble carbohydrates (glucose, fructose, and sucrose) of fruits from treated plants doubled concerning untreated plants.

Drought stress was also addressed by Kuvuran and Kuvuran (2019) in guar plants. There was a significant increase in antioxidant activity (total phenolic and flavonoid contents, superoxide dismutase, catalase, ascorbate peroxidase, and glutathione reductase), and reduction of oxidative stress (malondialdehyde) in guar plants treated with *Chlorella vulgaris*. Furthermore, the application of *A. platensis* in grape berries allowed the stomata to be open under water-stress conditions, without negatively impacting the water potential of vines, and promoted an increase in berry weight in both optimal and stress conditions. In addition, improved berry composition (sugar content) was also observed in treated vines under drought conditions (Salvi et al., 2020).

## Biocontrol effect

Pathogen organisms, such as insects, nematodes, bacteria, and fungi, strongly affect agricultural productivity. Some plants have their defense and resistance mechanisms, which include the regulation of signaling pathways, gene expression, and induction/inhibition of specific metabolic pathways, to produce secondary metabolites with

antioxidant and antimicrobial activities. However, the application of external protection agents is crucial to achieve the ambitious target productivity, to fulfill the growing food demands. Considering the wide variety of bioactive compounds that can be found in microalgae and cyanobacteria, the use of these microorganisms (or their extracts) can promote adequate crops protection against these biotic factors. Although the biocontrol effect was not so widely studied as the biofertilizer or biostimulant effect (Figure 4A), Table 6 provides published examples of the application of microalgae and cyanobacteria (biomass and/or extracts), on the biological control and protection of crops.

The fungicidal effect of microalgae against plant pathogens has been most widely evidenced than the antimicrobial effect. Only 3 recent studies from the last 2 years addressed the *in vitro* antibacterial effect of microalgae, or cyanobacteria, against plant pathogens (Al dayel et al., 2020; Bao et al., 2021; Ranglová et al., 2021). The potential of *Chlorella vulgaris* extracts as antibacterial was studied by Al dayel et al. (2020) and Ranglová et al. (2021), while Bao et al. (2021) evaluated the same effect from cyanobacterium *Anabaena variabilis*. Al dayel et al. (2020) obtained inhibition zones for *Escherichia coli*, *Streptococcus* sp., *Bacillus* sp., and *Staphylococcus aureus*, while Ranglová et al. (2021) showed that *C. vulgaris* grown in the synthetic medium were only active against *Clavibacter michiganensis*. *A. variabilis* extracts resulted in inhibition rates of 75.3 and 83.6% against *Xanthomonas oryzae* and *Rhizoctonia solani*, respectively, in *in vitro* trials (Bao et al., 2021). Moreover, *in vivo* trials were done by Bao et al. (2021) in rice fields infected with *Rhizoctonia solani*. The seedlings treated with *A. variabilis* were significantly longer than the control, achieving a disease control efficacy of 62.3%.

Kim (2006) evaluated different cyanobacteria isolates and Prasanna et al. (2008) evaluated several *Anabaena* isolates for biocidal activity against a set of phytopathogenic fungi. Nine cyanobacteria showed promising results. *Nostoc commune* and *Oscillatoria tenuis* showed strong antifungal activity against *Phytophthora capsica* (Kim, 2006). Among the *Anabaena* isolates, more than half showed inhibition zones of varying diameter against one or more fungi. The strains were characterized in terms of hydrolytic enzymes, proteins, and IAA (Indole-3-acetic acid), revealing a positive correlation between the presence of these metabolites and the fungicidal activity (Prasanna et al., 2008). Abdel-Hafez et al. (2015) also tested several cyanobacteria species and obtained a lower growth of *Alternaria porri*, using *Nostoc muscorum* and *Oscillatoria* sp. (20.4 and 36.3%, respectively). The culture filtrates of both species also contained high concentrations of phenolics compounds and alkaloids. Furthermore, their application in greenhouse conditions allowed for a reduction of 55.1-66.5% in the severity of purple blotch disease (Abdel-Hafez et al., 2015).

## Potential of wastewater-grown microalgae for agriculture

### Overview

Ecological management of manure on farms is vital to minimize losses of valuable plant nutrients and to prevent nutrient

TABLE 6 Impacts of microalgae and cyanobacteria on the enhancement of tolerance to biotic stress (biopesticide effect).

Biopesticide effect	Microalgae/ Cyanobacteria	Pest organisms	Reference
<b>Antibacterial</b>	<i>Chlorella vulgaris</i>	<i>Escherichia coli</i> , <i>Staphylococcus aureus</i> , <i>Streptococcus</i> sp., and <i>Bacillus</i> sp.	(Al dayel et al., 2020)
	<i>Anabaena variabilis</i>	<i>Xanthomonas oryzae</i>	(Bao et al., 2021)
<b>Fungicidal</b>	<i>Anabaena variabilis</i> <i>Anabaena minutissima</i>	<i>Rhizoctonia solani</i>	(Bao et al., 2021; Righini et al., 2021)
	<i>Anabaena</i> spp. <i>Calothrix</i> sp.	<i>Macrophomina phaseolina</i>	(Rana et al., 2012; Triveni et al., 2015)
	<i>Anabaena</i> spp.	<i>Macrophomina phaseolina</i> , <i>Fusarium moniliforme</i> , <i>Alternaria solani</i> , <i>Pythium debaryanum</i> , <i>Aspergillus candida</i> , <i>Drechslera oryzae</i> , <i>Fusarium solani</i>	(Prasanna et al., 2008)
	<i>Calothrix elenkinii</i>	<i>Pythium debaryanum</i> .	(Manjunath et al., 2010; Natarajan et al., 2012)
	<i>Chlorella fusca</i>	<i>Fusarium oxysporum</i> f. sp. <i>Fragariae</i>	(Kim et al., 2020)
	<i>Anabaena laxa</i> <i>Anabaena variabilis</i>	<i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i>	(Prasanna et al., 2013b)
	<i>Anabaena variabilis</i> <i>Anabaena oscillarioides</i>	Fungal consortium ( <i>Pythium debaryanum</i> , <i>Fusarium oxysporum lycopersici</i> , <i>Fusarium moniliforme</i> , and <i>Rhizoctonia solani</i> )	(Chaudhary et al., 2012)
	<i>Calothrix elenkinii</i> <i>Anabaena torulosa</i> <i>Anabaena laxa</i>	<i>Rhizoctonia</i>	(Prasanna et al., 2016b)
	<i>Anabaena oryzae</i> <i>Arthrospira</i> sp. <i>Nostoc minutum</i> <i>Nostoc muscorum</i> <i>Oscillatoria</i> sp.	<i>Alternaria porri</i>	(Abdel-Hafez et al., 2015)
	<i>Anabaena solitaria</i> <i>Anabaena</i> sp. <i>Calothrix brevissima</i> <i>Nostoc commune</i> <i>Nostoc muscorum</i> <i>Nodularia</i> sp. <i>Oscillatoria angustissima</i> <i>Oscillatoria tenuis</i>	<i>Alternaria alternata</i> , <i>Botrytis cineria</i> , <i>Colletotrichum gloeosporioides</i> , <i>Fusarium oxysporum</i> , <i>Phytophthora capsici</i> , <i>Pythium ultimum</i> , and <i>Rhizopus stolonifera</i>	(Kim, 2006)
<b>Anti-parasite</b>	<i>Oscillatoria chlorina</i>	<i>Meloidogyne arenaria</i>	(Khan et al., 2007)
<b>Insecticidal</b>	<i>Parachlorella kessleri</i> <i>Nostoc carneum</i>	<i>Spodoptera littoralis</i>	(Saber et al., 2018)

contamination of the surrounding watershed and soils. During the storage and land application of manure effluents, large amounts of N are lost to the atmosphere due to the volatilization of ammonia or nitrate runoff by rain to water streams and deep waters. An alternative to the land spreading of manure is to grow microalgae, where the pollutant from the effluents turns into nutrients from the microalgae point of view, converting them into biomass (Khan et al., 2019). Using microalgae for wastewater treatment (WWT) can provide a dual role, where bioremediation of wastewater occurs with the simultaneous production of valuable biomass, rich in essential nutrients and other bioactive compounds, which are beneficial for optimal seed germination and plant growth (amongst other applications). When microalgae are used for wastewater treatment, they are produced using open ponds due to their capacity to process large volumes of wastewater at lower

construction and operation costs, and easier scale-up. Acien et al. (2017) estimated the overall costs of microalgae-based WWT to be 30% lower than conventional activated sludge treatment, without even taking into account the revenues from potential commercial agricultural products.

Given the microalgal potential for WWT, there has been an increased interest in the use and research of these organisms, especially regarding the treatment of WW originating from livestock (i.e., piggeries, poultrys, aquaculture), humans (i.e., urban), industrial ones (i.e. breweries, wineries) and others such as foods and surface waters. Furthermore, with the scope of transforming WWTs into more circular processes, the biomass that is obtained is also being studied for application in agriculture (Table 7). The WW source will influence the biomass composition and, thus, the quality of the final product. The percentage of N and



TABLE 7 Summary of results for microalgae grown in wastewater and used for agricultural applications.

Wastewater	Culture medium	Microalgae	Agricultural application	References
Surface water	Eutrophic freshwater of three wastewater treatment facilities	Microalgae-cyanobacteria consortia	Composted MB sample (11.7, 21.6, 37.6% w/w) Improved signaling plant response to abiotic stress, plant growth, and biomass compared to control	(Ahn et al., 2020)
Urban	Centrate	<i>Chlorella vulgaris</i>	Freeze-dried microalga diluted with water (0.5 and 2 g/L) No significant effect on GI of cress seeds; No biopesticide activity	(Ranglová et al., 2021)
	75% of domestic wastewater	<i>Chlorella</i> sp. and <i>Scenedesmus</i> sp.	Deoiled microalgal biomass combined with inorganic fertilizer Improvement of tomato yield and nutrient content	(Silambarasan et al., 2021)
	Secondary wastewater	<i>Scenedesmus</i> sp.	Solid state formulations (2-9%) DW increase on ryegrass (220%) and barley plants (180%)	(González et al., 2020)
	6.5% (v/v) human urine	<i>Chlorella vulgaris</i>	Foliar spray of microalgal extracts (20-100%) Seeds treated with 40-60% microalgal extracts showed a faster plant growth rate	(Supraja et al., 2020)
		<i>Scenedesmus</i> sp.	Deoiled microalgal biomass as a single source or combined application with chemical fertilizer or vermicompost	(Nayak et al., 2019)
	Sewage	Microalgae-cyanobacteria consortia	Combined with 75% N and full-dose PK Higher NPK content in soil and plant; Increased plant weight compared to control	(Renuka et al., 2016)
Brewery	Anaerobically digested wastewater	<i>Tetrademus obliquus</i>	Microalga diluted with water (12.5% v/v) GI increase of 143% for barley seeds and 100% for wheat seeds Microalga extracts (0.1 and 0.5 g/L) 40% increase in GI of cress seeds using 0.1 g/L of untreated microalga	(Ferreira et al., 2019; Navarro-López et al., 2020b)
Dairy		<i>Chlorella pyrenoidosa</i>	Dried algal beads (20g) Increased root and shoot lengths of rice plants	(Yadavalli and Heggers, 2013)
Piggery	5% pig slurry	<i>Tetrademus obliquus</i> <i>Chlorella vulgaris</i> <i>Chlorella protothecoides</i> <i>Synechocystis</i> sp.	Microalgal cultures (0.5 g/L) Improved GI and root length of various seeds	(Ferreira et al., 2021)
		<i>Tetrademus obliquus</i>	GI increased by 40-45% for wheat seeds treated with 100-bar HPH microalga Non-treated <i>T. obliquus</i> increased up to 40% in root and shoot length Filtrate of the initial culture promoted shoot development (21%) compared to water	(Ferreira et al., 2022)
	10% pig manure	<i>Scenedesmus</i> sp.	Microalga extracts (0.1, 0.5, and 1 g/L) 12% increase in GI of cress seeds using 0.1 g/L of extracted microalga	(Navarro-López et al., 2020a)
Poultry	Water cooking of the poultry's remains from a slaughterhouse	<i>Chlorella vulgaris</i> <i>Tetrademus obliquus</i>	<i>C. vulgaris</i> (concentration) increased GI in 147% of wheat	(Viegas et al., 2021c)
Cattle	10% cattle manure pre-treated with ash	<i>Chlorella protothecoides</i> <i>Tetrademus obliquus</i>	Increments in the GI were 177% for wheat with <i>Chlorella protothecoides</i> and 34% for watercress with <i>Tetrademus obliquus</i>	(Viegas et al., 2021b)
	Anaerobically digested dairy manure		Seedlings grown in algae-amended potting mixes were equivalent to those grown with comparable levels of fertilizer-amended potting mixes with respect to seedling dry weight and nutrient content	(Mulbry et al., 2005)
Aquaculture	Brown crabs ( <i>Cancer pagurus</i> ) effluent	<i>Chlorella vulgaris</i> <i>Tetrademus obliquus</i>	Enhancements in GI for <i>C. vulgaris</i> (175%) and <i>T. obliquus</i> (48%) in watercress Enhancements in GI for <i>C. vulgaris</i> (84%) and <i>T. obliquus</i> (98%) in wheat	(Viegas et al., 2021a)
	Catfish water enriched with 30 g/L NaCl	<i>Arthrospira platensis</i>	Microalgal cultures (2-10 g/L) Improved seed germination and plant growth	(Wuang et al., 2016)

(Continued)

TABLE 7 Continued

Wastewater	Culture medium	Microalgae	Agricultural application	References
	Pikeperch recirculating freshwater	<i>Ulothrix</i> sp. and <i>Klebsormidium</i> sp.	Pasteurized dried microbial bacterial flocs Increase of NPK availability in soil; Lower fruit yield, but improved fruit quality (increase of 20% on DW, 23% in sugar content, and 70% on carotenoids)	(Coppens et al., 2016)
Food waste	Anaerobically digested wastewater	<i>Chlorella</i> sp. and <i>Scenedesmus</i> sp.	Dried biomass (adjusted to 100 kg N/ha) Better N retention	(Zarezadeh et al., 2019)
Meat processing	Water discarded from the sausage cooking and cooling tanks Cod desalting tanks and washing waters	<i>Chlorella vulgaris</i> , <i>Chlorolobium</i> sp., <i>Ankistrodesmus</i> sp., <i>Scenedesmus acutus</i> , <i>Oscillatoria</i> sp.	12% microalgal biomass added to fertilizer had a positive influence on plant development and its P recovery capacity	(Castro et al., 2017; de Souza et al., 2019; Castro et al., 2020)

GI, Germination Index (%); DW, Dry weight; HPH, High pressure homogenization.

P recovered by the biomass will depend largely on the initial concentration in the WW. Wastewaters generally have a low content of P, and its limitation has also been demonstrated to lead to an increased content of antioxidants (e.g., carotenoids, ascorbic acid, and tocopherols) (Gauthier et al., 2020). The presence of heavy metals, which can be accumulated by microalgae, also needs to be considered for the safety of agricultural microalgae-derived products (Ronga et al., 2019). The exposure of microalgae to heavy metals can trigger the synthesis of valuable compounds such as ascorbate peroxidase, catalase, superoxide dismutase, or ascorbate (Gauthier et al., 2020).

## Studies on the use of microalgal biomass in agriculture

The effect of unicellular and filamentous microalgal-cyanobacterial consortia, grown in urban wastewater, on the soil composition and grain quality of a wheat crop was evaluated (Renuka et al., 2016). Besides an increase in micronutrient availability in the soil (mainly zinc, iron, copper, and manganese), the studied consortia also enhanced their organic carbon content. This improvement in soil quality resulted in increased grain yield and higher product quality, as the obtained grains presented an improved nutritional composition. Silambarasan et al. (2021) studied the urban (at 75%) wastewater treatment capacity of a microalgal consortium composed of *Chlorella* sp. and *Scenedesmus* sp. The following use of deoiled microalgal biomass, combined with inorganic fertilizer, improved the tomato plant yield by 174%, as well as nutrient contents regarding nitrate (61%), phosphate (179%), potassium (71%), calcium (38%), magnesium (26%) and iron (11%). In another study, González et al. (2020) evaluated the formulation of microalgal biofertilizers from *Scenedesmus* sp. biomass obtained after the treatment of secondary urban wastewater. The application of solid-state formulations (2% microalgae) resulted in a DW increase of 220% and 180% for ryegrass and barley plants, respectively. Additionally, the authors observed an inhibitor effect on plant growth, for formulations that contained more than 2% microalgae. In another study, Ranglová et al. (2021) compared the biostimulant potential of two *C. vulgaris* cultures, grown in different media, inorganic BG11 medium, and centrate from municipal WW, by evaluating the

germination index of cress seeds, the auxin-like activity in mung bean and cytokinin-like activity in wheat growth tests. The biostimulant activity related to seed germination was found in the biomass cultivated in synthetic medium, but not for the one grown in WW, which could be related to the presence of inhibitory substances. The highest activity was obtained for *C. vulgaris* biomass harvested in the morning, at the lowest extract concentration. Both auxin-cytokinin-like activities were found in 2.0 and 3.0 g DW/L of *C. vulgaris* biomass harvested in the afternoon, which was equivalent to 0.5 mg DW/L of indole-3-butyric acid (auxin) and 0.3 mg DW/L of kinetin (cytokinin).

The research on the agricultural use of microalgal biomass applied in the treatment of animal-farming wastewater has also been gaining momentum. For instance, Ferreira et al. (2021) were able to improve the GI and root length of various plant seeds (tomato, watercress, cucumber, soybean, wheat, and barley seeds), after applying the biomass of a microalgal consortium composed of *Tetradesmus obliquus*, *Chlorella protothecoides*, *Chlorella vulgaris*, and the cyanobacterium *Synechocystis* sp., at 0.5g/L. In addition, Ferreira et al. (2022) studied the effect of *Tetradesmus obliquus* grown in 5% piggery effluent on wheat seed germination. The authors obtained three main conclusions: the use of biomass homogenized by high pressure (HPH) at 100-bar, or the use of non-treated biomass resulted in similar improvements of GI, up to 40-45% and 40%, respectively; and the use of filtrate from the initial culture also promoted shoot development up to 21%. These results demonstrate that all fractions of the microalgae culture have the potential to be utilized, including the supernatant as an enriched (in nutrients and bioactive compounds) irrigation solution and the microalga biomass in other high-value applications, as previously acknowledged by Kholssi et al. (2019). On another hand, Coppens et al. (2016) evaluated the application of microalgal-bacterial flocs (*Ulothrix* sp. and *Klebsormidium* sp.) in the treatment of brown crab's aquaculture wastewater, as well as the application of *Nannochloropsis oculata* on soil composition and fruit development in tomato plants. The microalgae biomass increased the availability of NPK in the soil, resulting in a plant growth comparable to the organic fertilizer treatment. Although a lower tomato yield was achieved than the treatment using inorganic fertilizer, the fruit quality was highly superior in terms of sugar and carotenoid contents.

In another study, Viegas et al. (2021a) analyzed the biostimulant effect of *Chlorella vulgaris* and *Tetradescmus obliquus*, used in the treatment of raw wastewater from brown crab aquaculture, on the germination of wheat and watercress seeds. The promising results obtained represented a GI improvement of 175 and 48% in watercress, and 84 and 98% in wheat, treated with *C. vulgaris* and *T. obliquus*, respectively.

Likewise, the agricultural potential of microalgal biomass obtained from the treatments of other wastewaters is also being explored. For example, Zarezadeh et al. (2019) improved the N retention in a common pasture of ryegrass, after the application of dried biomass from *Chlorella* sp. and *Scenedesmus* sp. grown in anaerobically digested wastewater from food waste. The authors observed a slow release of nutrients which ultimately impacted the N cycling through the modulation of N mineralization and N pathways. In addition, de Souza et al. (2019) performed a life cycle analysis of the application on soil, of 1 kg of N from microalgal biomass grown in food waste effluent. The authors suggested the alternative uses of photovoltaic energy and nitrogen-rich effluents, as well as the consideration of environmental compensation for the treatment of effluent, to improve the economic and environmental viability of the use of microalgal N biofertilizer. In another case, Ferreira et al. (2019) were able to increase the germination percentage of barley and wheat seeds, by 143 and 100%, respectively, with the application of *Tetradescmus obliquus* used in the treatment of anaerobically digested wastewater from a brewery. Finally, Ahn et al. (2020) evaluated the effects of the use of biomass of microalgae-cyanobacterium consortia, obtained in eutrophic freshwater from three wastewater treatment facilities, on *Perilla* sp. plants. Positive results were registered, with the improvement of plant signaling to abiotic stress, plant growth, and biomass, thus setting forth one more high potential application of microalgal biomass obtained with the treatment of wastewaters.

## Conclusions

The practical applications of microalgae and cyanobacteria in agriculture are still very limited since the focus on these microorganisms has been on their functional activities in food and feed (as nutraceuticals and additives), pharmaceutical, and cosmetic products. However, their interest has been increasing given their potential role in enriching soil quality and plant nutrition, stimulating plant growth, and expanding plant tolerance to biotic and abiotic stress.

Although cyanobacteria, commonly known as blue-green algae, have already been recognized for their fertilizing role on the soil and rhizosphere, the research on the application of microalgae for biostimulants and biopesticides has drastically increased in the last two years. Most works have shown evidence of microalgae improving plant productivity and resistance to adverse conditions, thus supporting them as a viable biological-based alternative to chemical fertilizers, pesticides, and growth stimulants. Moreover,

some studies highlight the added benefit of microalgae in enhancing the biometric and nutritional quality of leaves and fruits.

Despite the recognized benefits, the mode of action of microalgal compounds on plant response mechanisms is still not fully understood, given the high diversity of species and their products. Hence, metabolomics' studies focusing on the effect of each metabolite/biomass on crops are essential to identify the most adequate for agricultural activities. Furthermore, only a small set of studies have been developed using microalgae grown in wastewaters, although with promising results. Nevertheless, until now no data have been reported on the specific composition of an algal culture medium that could help to understand its positive biofertilizing effect; therefore, the characterization of compounds with biofertilizing capability and enzymatic activity profiles in the extracellular medium of algal cultures also needs to be investigated.

The use of microalgae in sustainable agriculture is expected to be one of the most important applications of microalgal biomass in short term, especially for biomass cultivated in wastewater. This will not only bring benefits from an environmental perspective, concerning the recovery of nutrients and recycling of water in wastewater treatments and agriculture but will also improve the economic feasibility of microalgal production and downstream processing.

With increasing climate change, global warming, water scarcity, degradation of soils, and emerging crisis such as the war in Ukraine, innovations will be needed to enhance and protect crops throughout the world. The challenge to produce more food with limited resources makes microalgae a suitable alternative for enhancing and protecting agricultural production and delivering economic and environmental benefits to farmers and algae producers.

## Author contributions

Conceptualization, AF. Writing—original draft preparation, AF. Writing—review and editing, AF, CB, CM-D-S, FA-F and LG. Visualization, AF, and CB. Data curation, CB. Supervision, CM-D-S, FA-F and LG. Project administration, LG. Funding acquisition, LG. All authors contributed to the article and approved the submitted version.

## Funding

This work was supported by the project ALGAVALOR - Lisboa-01-0247-FEDER-035234, supported by Operational Programme for Competitiveness and Internationalization (COMPETE2020), by Lisbon Portugal Regional Operational Programme (Lisboa 2020) and by Algarve Regional Operational Programme (Algarve 2020) under the Portugal 2020 Partnership Agreement, through the European Regional Development Fund (ERDF); Biomass and Bioenergy Research Infrastructure (BBRI)-LISBOA-01-0145-FEDER-022059, supported by Operational Programme for Competitiveness and Internationalization

(PORTUGAL2020), by Lisbon Portugal Regional Operational Programme (Lisboa 2020) and by North Portugal Regional Operational Programme (Norte 2020) under the Portugal 2020 Partnership Agreement, through the European Regional Development Fund (ERDF); project Performalgae, from the European Union's Horizon 2020 research and innovation programme (grant agreement n° ALG-01-0247-FEDER-069961); Red CYTED P319RT0025 - RENUWAL - Red Iberoamericana para el Tratamiento de Efluentes con Microalgas; Bilateral Portugal-India DRI/India/0609/2020 Project WCAlgaeKIT<sup>+</sup> - Combination of Vertical Wetlands, Microalgae Photobioreactor, and Microbial Fuel Cell (KIT) for wastewater treatment in small pig production farms. AF and CB are pleased to acknowledge their PhD grants SFRH/BD/144122/2019 and FCT 2022.14298.BDANA, respectively, awarded by Fundação para a Ciência e Tecnologia.

## References

- Abdel-Hafez, S. I. I., Abo-Elyousr, K. A. M., and Abdel-Rahim, I. R. (2015). Fungicidal activity of extracellular products of cyanobacteria against *Alternaria porri*. *Eur. J. Phycol.* 50, 239–245. doi: 10.1080/09670262.2015.1028105
- Abdel-Raouf, N. (2012). Agricultural importance of algae. *Afr. J. Biotechnol.* 11, 11648–11658. doi: 10.5897/AJB11.3983
- Acea, M. J., Diz, N., and Prieto-Fernández, A. (2001). Microbial populations in heated soils inoculated with cyanobacteria. *Biol. Fertil. Soils.* 33, 118–125. doi: 10.1007/s003740000298
- Acien, F. G., Molina, E., Fernández-Sevilla, J. M., Barbosa, M., Gouveia, L., Sepúlveda, C., et al. (2017). “20 - Economics of microalgae production,” in *Microalgae-Based Biofuels and Bioproducts: From Feedstock Cultivation to End-products*. Eds. C. Gonzalez-Fernandez and R. Muñoz (Duxford, UK: Woodhead Publishing), 485–503.
- Ahn, C. H., Lee, S., Park, J. R., Hwang, T.-M., and Joo, J. C. (2020). Harvested microalgal biomass from different water treatment facilities—its characteristics and potential use as renewable sources of plant biostimulation. *Agronomy* 10, 1882. doi: 10.3390/agronomy10121882
- Akgül, F. (2019). Effect of *Spirulina platensis* (Gomont) geitler extract on seed germination of wheat and barley. *Alinteri. Zirai. Bilim. Derg.* 34, 148–153. doi: 10.28955/alinterizbd.639000
- Al dayel, M. F., El Semary, N. A., Al Amer, K., and Al Ali, K. M. (2020). Investigating the applications of *Chlorella vulgaris* in agriculture and nanosilver production. *J. Environ. Biol.* 41, 1099–1104. doi: 10.22438/jeb/41/5/MRN-1395
- Atzori, G., Nissim, W. G., Rodolfi, L., Niccolai, A., Biondi, N., Mancuso, S., et al. (2020). Algae and bioguanos as promising source of organic fertilizers. *J. Appl. Phycol.* 32, 3971–3981. doi: 10.1007/s10811-020-02261-7
- Bao, J., Zhuo, C., Zhang, D., Li, Y., Hu, F., Li, H., et al. (2021). Potential applicability of a cyanobacterium as a biofertilizer and biopesticide in rice fields. *Plant Soil* 463, 1–16. doi: 10.1007/s11104-021-04899-9
- Barone, V., Baglieri, A., Stevanato, P., Broccanello, C., Bertoldo, G., Bertaggia, M., et al. (2018). Root morphological and molecular responses induced by microalgae extracts in sugar beet (*Beta vulgaris* L.). *J. Appl. Phycol.* 30, 1061–1071. doi: 10.1007/s10811-017-1283-3
- Barone, V., Puglisi, I., Fragalà, F., Lo Piero, A. R., Giuffrida, F., and Baglieri, A. (2019a). Novel bioprocess for the cultivation of microalgae in hydroponic growing system of tomato plants. *J. Appl. Phycol.* 31, 465–470. doi: 10.1007/s10811-018-1518-y
- Barone, V., Puglisi, I., Fragalà, F., Stevanato, P., and Baglieri, A. (2019b). Effect of living cells of microalgae or their extracts on soil enzyme activities. *Arch. Agron. Soil Sci.* 65, 712–726. doi: 10.1080/03650340.2018.1521513
- Barsanti, L., Coltelli, P., and Gualtieri, P. (2019). Paramylon treatment improves quality profile and drought resistance in *Solanum lycopersicum* L. cv. micro-tom. *Agronomy* 9, 394. doi: 10.3390/agronomy9070394
- Baweja, P., Kumar, S., and Kumar, G. (2019). “Organic fertilizer from algae: a novel approach towards sustainable agriculture,” *Biofertilizers for Sustainable Agriculture and Environment. Soil Biology*. Eds. B. Giri, R. Prasad, Q. S. Wu and A. Varma (Cham, Switzerland: Springer), 353–370.
- Bayona-Morcillo, P. J., Plaza, B. M., Gómez-Serrano, C., Rojas, E., and Jiménez-Becker, S. (2020). Effect of the foliar application of cyanobacterial hydrolysate (*Arthrospira platensis*) on the growth of *Petunia x hybrida* under salinity conditions. *J. Appl. Phycol.* 32, 4003–4011. doi: 10.1007/s10811-020-02192-3
- Boopathi, T., Balamurugan, V., Gopinath, S., and Sundararaman, M. (2013). Characterization of IAA production by the mangrove cyanobacterium *Phormidium* sp. MI405019 and its influence on tobacco seed germination and organogenesis. *J. Plant Growth Regul.* 32, 758–766. doi: 10.1007/s00344-013-9342-8
- Buenaventura, M. K. P., and Barrientos, D. S. (2019). Response of *Oryza sativa* CL1 (Basmati 370) to *Nostoc commune* vaucl. as fertilizer supplement. *Mindanao. J. Sci. Technol.* 17, 242–256.
- Bulgari, R., Cocetta, G., Trivellini, A., Vernieri, P., and Ferrante, A. (2015). Biostimulants and crop responses: a review. *Biol. Agric. Hortic.* 31, 1–17. doi: 10.1080/01448765.2014.964649
- Calvo, P., Nelson, L., and Kloepper, J. W. (2014). Agricultural uses of plant biostimulants. *Plant Soil* 383, 3–41. doi: 10.1007/s11104-014-2131-8
- Carvajal-Muñoz, J. S., and Carmona-García, C. E. (2012). Benefits and limitations of biofertilization in agricultural practices. *Livest. Res. Rural Dev.* 24.
- Castro, J., de, S., Calijuri, M. L., Assemany, P. P., Cecon, P. R., de Assis, I. R., et al. (2017). Microalgal biofilm in soil: greenhouse gas emissions, ammonia volatilization and plant growth. *Sci. Total. Environ.* 574, 1640–1648. doi: 10.1016/j.scitotenv.2016.08.205
- Castro, J., de, S., Calijuri, M. L., Mattiello, E. M., Ribeiro, V. J., and Assemany, P. P. (2020). Algal biomass from wastewater: soil phosphorus bioavailability and plants productivity. *Sci. Total. Environ.* 711, 135088. doi: 10.1016/j.scitotenv.2019.135088
- Chanda, M., Merghoub, N., and EL Arroussi, H. (2019). Microalgae polysaccharides: the new sustainable bioactive products for the development of plant bio-stimulants? *World J. Microbiol. Biotechnol.* 35, 1–10. doi: 10.1007/s11274-019-2745-3
- Chatterjee, A., Singh, S., Agrawal, C., Yadav, S., Rai, R., and Rai, L. C. (2017). “Role of algae as a biofertilizer,” *Algal Green Chemistry: Recent Progress in Biotechnology*. Eds. R. P. Rastogi, A. Pandey and D. Madamwar (Amsterdam, Netherlands: Elsevier), 189–200.
- Chaudhary, V., Prasanna, R., Nain, L., Dubey, S. C., Gupta, V., Singh, R., et al. (2012). Bioefficacy of novel cyanobacteria-amended formulations in suppressing damping off disease in tomato seedlings. *World J. Microbiol. Biotechnol.* 28, 3301–3310. doi: 10.1007/S11274-012-1141-Z
- Chew, P., and Soccio, M. (2016). *Asia-Pacific: agricultural perspectives*. Available at: <https://economics.rabobank.com/publications/2016/february/asia-pacific-agricultural-perspectives/> (Accessed July 6, 2021).
- Chiaiese, P., Corrado, G., Colla, G., Kyriacou, M. C., and Roupheal, Y. (2018). Renewable sources of plant biostimulation: microalgae as a sustainable means to improve crop performance. *Front. Plant Sci.* 871. doi: 10.3389/fpls.2018.01782
- Chua, A., Sherwood, O. L., Fitzhenry, L., Ng, C. K. Y., McCabe, P. F., and Daly, C. T. (2020). Cyanobacteria-derived proline increases stress tolerance in arabidopsis thaliana root hairs by suppressing programmed cell death. *Front. Plant Sci.* 11. doi: 10.3389/fpls.2020.490075
- Colla, G., and Roupheal, Y. (2015). Biostimulants in horticulture. *Sci. Hortic. (Amsterdam)*. 196, 1–2. doi: 10.1016/j.scienta.2015.10.044
- Coppens, J., Grunert, O., Van Den Hende, S., Vanhoutte, I., Boon, N., Haesaert, G., et al. (2016). The use of microalgae as a high-value organic slow-release fertilizer results in tomatoes with increased carotenoid and sugar levels. *J. Appl. Phycol.* 28, 2367–2377. doi: 10.1007/s10811-015-0775-2
- Corbellini, J. R., Ribas, L. L. F., de Maia, F. R., Corrêa, D., de, O., Nosedá, M. D., et al. (2020). Effect of microalgae *Messastrium gracile* and *Chlorella vulgaris* on the *in vitro*

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.



- propagation of orchid *Cattleya labiata*. *J. Appl. Phycol.* 32, 4013–4027. doi: 10.1007/s10811-020-02251-9
- Costa, J. A. V., Freitas, B. C. B., Cruz, C. G., Silveira, J., and Moraes, M. G. (2019). Potential of microalgae as biopesticides to contribute to sustainable agriculture and environmental development. *J. Environ. Sci. Heal. - Part B. Pestic. Food Contam. Agric. Wastes*. 54, 366–375. doi: 10.1080/03601234.2019.1571366
- Decesaro, A., Rampel, A., Machado, T. S., Thomé, A., Reddy, K., Margarites, A. C., et al. (2016). Bioremediation of soil contaminated with diesel and biodiesel fuel using biostimulation with microalgae biomass. *J. Environ. Eng.* 143, 04016091. doi: 10.1061/(ASCE)EE.1943-7870.0001165
- Deepika, P., and MubarakAli, D. (2020). Production and assessment of microalgal liquid fertilizer for the enhanced growth of four crop plants. *Biocatal. Agric. Biotechnol.* 28, 101701. doi: 10.1016/j.bcab.2020.101701
- de Souza, M. H. B., Calijuri, M. L., Assemany, P. P., Castro, J., de, S., and de Oliveira, A. C. M. (2019). Soil application of microalgae for nitrogen recovery: a life-cycle approach. *J. Clean. Prod.* 211, 342–349. doi: 10.1016/j.jclepro.2018.11.097
- Dias, G. A., Rocha, R. H. C., Araújo, J. L., De Lima, J. F., and Guedes, W. A. (2016). Growth, yield, and postharvest quality in eggplant produced under different foliar fertilizer (*Spirulina platensis*) treatments. *Semin. Agrar.* 37, 3893–3902. doi: 10.5433/1679-0359.2016v37n6p3893
- Dineshkumar, R., Duraimurugan, M., Sharmiladevi, N., Lakshmi, L. P., Rasheeq, A. A., Arumugam, A., et al. (2020a). Microalgal liquid biofertilizer and biostimulant effect on green gram (*Vigna radiata* L.) an experimental cultivation. *Biomass Convers. Biorefinery*. 12, 3007–3027. doi: 10.1007/s13399-020-00857-0
- Dineshkumar, R., Kumaravel, R., Gopalsamy, J., Sikder, M. N. A., and Sampathkumar, P. (2018). Microalgae as bio-fertilizers for rice growth and seed yield productivity. *Waste. Biomass Valorization*. 9, 793–800. doi: 10.1007/s12649-017-9873-5
- Dineshkumar, R., Subramanian, J., Arumugam, A., Ahamed Rasheeq, A., and Sampathkumar, P. (2020b). Exploring the microalgae biofertilizer effect on onion cultivation by field experiment. *Waste. Biomass Valorization*. 11, 77–87. doi: 10.1007/s12649-018-0466-8
- Dineshkumar, R., Subramanian, J., Gopalsamy, J., Jayasingam, P., Arumugam, A., Kannadasan, S., et al. (2019). The impact of using microalgae as biofertilizer in maize (*Zea mays* L.). *Waste. Biomass Valorization*. 10, 1101–1110. doi: 10.1007/s12649-017-0123-7
- Dineshkumar, R., Subramanian, J., and Sampathkumar, P. (2020c). Prospective of *Chlorella vulgaris* to augment growth and yield parameters along with superior seed qualities in black gram, *Vigna mungo* (L.). *Waste. Biomass Valorization*. 11, 1279–1287. doi: 10.1007/s12649-018-0465-9
- Do, T. C., Van, Tran, D. T., Le, T. G., and Nguyen, Q. T. (2020). Characterization of endogenous auxins and gibberellins produced by *Chlorella sorokiniana* TH01 under phototrophic and mixotrophic cultivation modes toward applications in microalgal biorefinery and crop research. *J. Chem.* 2020, 4910621. doi: 10.1155/2020/4910621
- Dommm, P. (2022). A fertilizer shortage, worsened by war in Ukraine, is driving up global food prices and scarcity (CNBC). Available at: <https://www.cnbc.com/2022/04/06/a-fertilizer-shortage-worsened-by-war-in-ukraine-is-driving-up-global-food-prices-and-scarcity.html> (Accessed July 4, 2022).
- Ekinici, K., Erdal, I., Uysal, Ö., Uysal, F. Ö., Tuncce, H., and Doğan, A. (2019). Anaerobic digestion of three microalgae biomasses and assessment of digestates as biofertilizer for plant growth. *Environ. Prog. Sustain. Energy* 38, 1–9. doi: 10.1002/ep.13024
- El Arroussi, H., Benhima, R., Elbaouchi, A., Sijlmasi, B., EL Mernissi, N., Aafsar, A., et al. (2018). *Dunaliella salina* exopolysaccharides: a promising biostimulant for salt stress tolerance in tomato (*Solanum lycopersicum*). *J. Appl. Phycol.* 30, 2929–2941. doi: 10.1007/s10811-017-1382-1
- El Arroussi, H., El Mernissi, N., Benhima, R., El Kadmiri, I. M., Bendaou, N., and Smouni, A. (2016). Microalgae polysaccharides a promising plant growth biostimulant. *J. Algal. Biomass Utiln.* 7, 55–63.
- El-Naggar, A. H., Osman, M. E. H. E.-S., and Gheda, S. F. (2005). Influence of the aqueous extracts of *Ulva lactuca* and *Chlorella kessleri* on growth and yield of *Vicia faba*. *Algol. Stud. für. Hydrobiol. Suppl.* Vol. 116, 213–229. doi: 10.1127/1864-1318/2005/0116-0213
- El Semary, N. A. H., Alouane, M. H. H., Nasr, O., Aldayel, M. F., Alhaweti, F. H., and Ahmed, F. (2020). Salinity stress mitigation using encapsulated biofertilizers for sustainable agriculture. *Sustainability* 12, 9218. doi: 10.3390/su12219218
- El-Sheekh, M. M., Zayed, M. A., Elmossel, F. K. A., and Hassan, R. S. A. (2018). Effect of cyanobacteria isolates on rice seeds germination in saline soil. *Baghdad. Sci. J.* 15, 16–21. doi: 10.21123/bsj.2018.15.1.0016
- Eurostat (2020). Agri-environmental indicator - consumption of pesticides. In: *Eurostat - stat. explain*. Available at: [https://ec.europa.eu/eurostat/statistics-explained/index.php/Agri-environmental\\_indicator\\_-\\_consumption\\_of\\_pesticides#Analysis\\_at\\_EU\\_and\\_country\\_level](https://ec.europa.eu/eurostat/statistics-explained/index.php/Agri-environmental_indicator_-_consumption_of_pesticides#Analysis_at_EU_and_country_level) (Accessed April 8, 2021).
- Faheed, F. A., and Fattah, Z. A.-E. (2008). Effect of *Chlorella vulgaris* as bio-fertilizer on growth parameters and metabolic aspects of lettuce plant. *J. Agric. Soc. Sci.* 4, 165–169.
- Farid, R., Mutale-joan, C., Redouane, B., Mernissi Najib, E., Abderahime, A., Laila, S., et al. (2019). Effect of microalgae polysaccharides on biochemical and metabolomics pathways related to plant defense in *Solanum lycopersicum*. *Appl. Biochem. Biotechnol.* 188, 225–240. doi: 10.1007/s12010-018-2916-y
- Ferreira, A., Figueiredo, D., Ferreira, F., Ribeiro, B., Reis, A., Silva, T. L., et al. (2022). Impact of high-pressure homogenization on the cell integrity of *tetrademus obliquus* and seed germination. *Molecules* 27, 2275. doi: 10.3390/MOLECULES27072275
- Ferreira, A., Melkonyan, L., Carapinha, S., Ribeiro, B., Figueiredo, D., Avetisova, G., et al. (2021). Biostimulant and biopesticide potential of microalgae growing in piggery wastewater. *Environ. Adv.* 4, 100062. doi: 10.1016/j.envadv.2021.100062
- Ferreira, A., Ribeiro, B., Ferreira, A. F., Tavares, M. L. A., Vlado, J., Vidović, S., et al. (2019). *Scenedesmus obliquus* microalga-based biorefinery – from brewery effluent to bioactive compounds, biofuels and biofertilizers – aiming at a circular bioeconomy. *Biofuels. Bioprod. Biorefining*. 13, 1169–1186. doi: 10.1002/bbb.2032
- Garcia-Gonzalez, J., and Sommerfeld, M. (2016). Biofertilizer and biostimulant properties of the microalga *Acutodesmus dimorphus*. *J. Appl. Phycol.* 28, 1051–1061. doi: 10.1007/s10811-015-0625-2
- Gauthier, M. R., Senhorinho, G. N. A., and Scott, J. A. (2020). Microalgae under environmental stress as a source of antioxidants. *Algal. Res.* 52, 102104. doi: 10.1016/J.ALGAL.2020.102104
- Gayathri, M., Shunmugam, S., Thajuddin, N., and Muralitharan, G. (2017). Phytohormones and free volatile fatty acids from cyanobacterial biomass wet extract (BWE) elicit plant growth promotion. *Algal. Res.* 26, 56–64. doi: 10.1016/j.algal.2017.06.022
- Gemin, L. G., Mógor, Á.F., De Oliveira Amatusi, J., and Mógor, G. (2019). Microalgae associated to humic acid as a novel biostimulant improving onion growth and yield. *Sci. Hortic. (Amsterdam)*. 256, 108560. doi: 10.1016/j.scienta.2019.108560
- Godlewska, K., Michalak, I., Pacyga, P., Baśladyńska, S., and Chojnacka, K. (2019). Potential applications of cyanobacteria: spirulina platensis filtrates and homogenates in agriculture. *World J. Microbiol. Biotechnol.* 35, 80. doi: 10.1007/s11274-019-2653-6
- Gonçalves, A. L. (2021). The use of microalgae and cyanobacteria in the improvement of agricultural practices: a review on their biofertilising, biostimulating and biopesticide roles. *Appl. Sci.* 11, 1–21. doi: 10.3390/app11020871
- González, I., Herrero, N., Siles, J. Á., Chica, A. F., Ángeles Martín, M., Izquierdo, C. G., et al. (2020). Wastewater nutrient recovery using twin-layer microalgae technology for biofertilizer production. *Water Sci. Technol.* 82, 1044–1061. doi: 10.2166/wst.2020.372
- Górka, B., Korzeniowska, K., Lipok, J., and Wiczorek, P. P. (2018). “The biomass of algae and algal extracts in agricultural production,” in *Algae biomass: characteristics and applications*. Eds. K. Chojnacka, P. Wiczorek, G. Schroeder and I. Michalak (Cham, Switzerland: Springer International Publishing) 8, 103–114.
- Górka, B., Lipok, J., and Wiczorek, P. P. (2015). “Biologically active organic compounds, especially plant promoters, in algae extracts and their potential application in plant cultivation,” in *Marine algae extracts* (Weinheim, Germany: Wiley-VCH Verlag GmbH & Co. KGaA), 659–680. doi: 10.1002/9783527679577.ch37
- Grzesik, M., and Romanowska-Duda, Z. (2014). Improvements in germination, growth, and metabolic activity of corn seedlings by grain conditioning and root application with cyanobacteria and microalgae. *Polish. J. Environ. Stud.* 23, 1147–1153.
- Grzesik, M., Romanowska-Duda, Z., and Kalaji, H. M. (2017). Effectiveness of cyanobacteria and green algae in enhancing the photosynthetic performance and growth of willow (*Salix viminalis* L.) plants under limited synthetic fertilizers application. *Photosynthetica* 55, 510–521. doi: 10.1007/s11099-017-0716-1
- Guzmán-Murillo, M. A., Ascencio, F., and Larrinaga-Mayoral, J. A. (2013). Germination and ROS detoxification in bell pepper (*Capsicum annuum* L.) under NaCl stress and treatment with microalgae extracts. *Protoplasma* 250, 33–42. doi: 10.1007/s00709-011-0369-z
- Hu, C., Liu, Y., Song, L., and Zhang, D. (2002). Effect of desert soil algae on the stabilization of fine sands. *J. Appl. Phycol.* 14, 281–292. doi: 10.1023/A:1021128530086
- Hussain, A., and Hasnain, S. (2011). Phytostimulation and biofertilization in wheat by cyanobacteria. *J. Ind. Microbiol. Biotechnol.* 38, 85–92. doi: 10.1007/s10295-010-0833-3
- Issa, O. M., Défarge, C., Le Bissonnais, Y., Marin, B., Duval, O., Bruand, A., et al. (2007). Effects of the inoculation of cyanobacteria on the microstructure and the structural stability of a tropical soil. *Plant Soil* 290, 209–219. doi: 10.1007/s11104-006-0060-6
- Karthikeyan, N., Prasanna, R., Nain, L., and Kaushik, B. D. (2007). Evaluating the potential of plant growth promoting cyanobacteria as inoculants for wheat. *Eur. J. Soil Biol.* 43, 23–30. doi: 10.1016/j.ejsobi.2006.11.001
- Khan, Z., Kim, Y. H., Kim, S. G., and Kim, H. W. (2007). Observations on the suppression of root-knot nematode (*Meloidogyne arenaria*) on tomato by incorporation of cyanobacterial powder (*Oscillatoria chlorina*) into potting field soil. *Bioresour. Technol.* 98, 69–73. doi: 10.1016/J.BIORTECH.2005.11.029
- Khan, S. A., Sharma, G. K., Malla, F. A., Kumar, A., Rashmi, and Gupta, N. (2019). Microalgae based biofertilizers: a biorefinery approach to phycoremediate wastewater and harvest biodiesel and manure. *J. Clean. Prod.* 211, 1412–1419. doi: 10.1016/j.jclepro.2018.11.281
- Kheirfam, H., Sadeghi, S. H., Homae, M., and Zarei Darki, B. (2017). Quality improvement of an erosion-prone soil through microbial enrichment. *Soil Tillage. Res.* 165, 230–238. doi: 10.1016/j.still.2016.08.021



- Kholssi, R., Marks, E. A. N., Miñón, J., Montero, O., Deboudi, A., and Rad, C. (2019). Biofertilizing effect of chlorella sorokiniana suspensions on wheat growth. *J. Plant Growth Regul.* 38, 644–649. doi: 10.1007/s00344-018-9879-7
- Kim, J.-D. (2006). Screening of cyanobacteria (Blue-green algae) from rice paddy soil for antifungal activity against plant pathogenic fungi. *Mycobiology* 34, 138. doi: 10.4489/MYCO.2006.34.3.138
- Kim, M. J., Shim, C. K., Ko, B. G., and Kim, J. (2020). Effect of the microalga chlorella fusca CHK0059 on strawberry PGPR and biological control of fusarium wilt disease in non-pesticide hydroponic strawberry cultivation. *J. Microbiol. Biotechnol.* 30, 708–716. doi: 10.4014/jmb.2001.01015
- Kopta, T., Pavliková, M., Šekara, A., Pokluda, R., and Maršálek, B. (2018). Effect of bacterial-algal biostimulant on the yield and internal quality of lettuce (*Lactuca sativa* L.) produced for spring and summer crop. *Not. Bot. Horti. Agrobot. Cluj-Napoca*. 46, 615–621. doi: 10.15835/nbha46211110
- Kusvuran, A., and Kusvuran, S. (2019). Using of microbial fertilizer as biostimulant alleviates damage from drought stress in guar (*Cyamopsis tetragonoloba* (L.) taub.) seedlings. *Int. Lett. Nat. Sci.* 76, 147–157. doi: 10.18052/www.scipress.com/ilns.76.147
- Liebig, N., Salaun, M., and Monnier, C. (2020). Development of standards and guidance documents for biostimulants approval under European fertilizer regulation (EU) 2019/1009. *Eurofins. Agros. Serv.* 2019–2021.
- Loganathan, B. G., Orsat, V., and Lefsrud, M. (2020). Utilizing the microalgal biomass of chlorella variabilis and scenedesmus obliquus produced from the treatment of synthetic dairy wastewater as a biofertilizer. *J. Plant Nutr.* 44, 1486–1497. doi: 10.1080/01904167.2020.1862191
- Lv, J., Liu, S., Feng, J., Liu, Q., Guo, J., Wang, L., et al. (2020). Effects of microalgal biomass as biofertilizer on the growth of cucumber and microbial communities in the cucumber rhizosphere. *Turk. J. Bot.* 44, 167–177. doi: 10.3906/bot-1906-1
- Manjunath, M., Kanchan, A., Ranjan, K., Venkatachalam, S., Prasanna, R., Ramakrishnan, B., et al. (2016). Beneficial cyanobacteria and eubacteria synergistically enhance bioavailability of soil nutrients and yield of okra. *Heliyon* 2, e00066. doi: 10.1016/J.HELIYON.2016.E00066
- Manjunath, M., Prasanna, R., Nain, L., Dureja, P., Singh, R., Kumar, A., et al. (2010). Biocontrol potential of cyanobacterial metabolites against damping off disease caused by *Pythium aphanidermatum* in solanaceous vegetables. *Arch. Phytopathol. Plant Prot.* 43, 666–677. doi: 10.1080/03235400802075815
- MarketsandMarkets (2020). *Biostimulants market by active ingredient, crop type, application method, form - global forecast 2025*. Available at: <https://www.marketsandmarkets.com/Market-Reports/biostimulant-market-1081.html> (Accessed September 7, 2020).
- Marks, E. A. N., Miñón, J., Pascual, A., Montero, O., Navas, L. M., and Rad, C. (2017). Application of a microalgal slurry to soil stimulates heterotrophic activity and promotes bacterial growth. *Sci. Total. Environ.* 605–606, 610–617. doi: 10.1016/j.scitotenv.2017.06.169
- Marks, E. A. N., Montero, O., and Rad, C. (2019). The biostimulating effects of viable microalgal cells applied to a calcareous soil: increases in bacterial biomass, phosphorus scavenging, and precipitation of carbonates. *Sci. Total. Environ.* 692, 784–790. doi: 10.1016/j.scitotenv.2019.07.289
- Mazhar, S., and Hasnain, S. (2011). Screening of native plant growth promoting cyanobacteria and their impact on triticum aestivum var. uqab 2000 growth. *Afr. J. Agric. Res.* 6, 3988–3993. doi: 10.5897/AJAR11.560
- Michalak, I., Chojnacka, K., Dmytryk, A., Wilk, R., Gramza, M., and Rój, E. (2016). Evaluation of supercritical extracts of algae as biostimulants of plant growth in field trials. *Front. Plant Sci.* 7. doi: 10.3389/fpls.2016.01591
- Mógor, Á.F., de Oliveira Amatusi, J., Mógor, G., and Bocchetti de Lara, G. (2018a). Bioactivity of cyanobacterial biomass related to amino acids induces growth and metabolic changes on seedlings and yield gains of organic red beet. *Am. J. Plant Sci.* 09, 966–978. doi: 10.4236/ajps.2018.95074
- Mógor, Á.F., Ördög, V., Lima, G. P. P., Molnár, Z., and Mógor, G. (2018b). Biostimulant properties of cyanobacterial hydrolysate related to polyamines. *J. Appl. Phycol.* 30, 453–460. doi: 10.1007/s10811-017-1242-z
- Mordor Intelligence (2020). *Europe Biostimulants market - growth, trends and forecasts, (2020-2025)*. Available at: <https://www.mordorintelligence.com/industry-reports/europe-biostimulants-market> (Accessed September 7, 2020).
- Mukherjee, C., Chowdhury, R., Sutradhar, T., Begam, M., Ghosh, S. M., Basak, S. K., et al. (2016). Parboiled rice effluent: a wastewater niche for microalgae and cyanobacteria with growth coupled to comprehensive remediation and phosphorus biofertilization. *Algal Res.* 19, 225–236. doi: 10.1016/j.algal.2016.09.009
- Mulbry, W., Westhead, E. K., Pizarro, C., and Sikora, L. (2005). Recycling of manure nutrients: use of algal biomass from dairy manure treatment as a slow release fertilizer. *Bioresour. Technol.* 96, 451–458. doi: 10.1016/j.biortech.2004.05.026
- Mutale-joan, C., Redouane, B., Najib, E., Yassine, K., Lyamlouli, K., Laila, S., et al. (2020). Screening of microalgae liquid extracts for their bio stimulant properties on plant growth, nutrient uptake and metabolite profile of solanum lycopersicum L. *Sci. Rep.* 10, 1–12. doi: 10.1038/s41598-020-59840-4
- Narain, D. (2020). *Transforming Indian agriculture: a policy framework t guide US-India partnership* (Washington DC). Available at: <http://www.fao.org/3/a/> (Accessed July 6, 2021).
- Natarajan, C., Prasanna, R., Gupta, V., Dureja, P., and Nain, L. (2012). Characterization of the fungicidal activity of calothrix elenkini using chemical methods and microscopy. *Appl. Biochem. Microbiol.* 48, 51–57. doi: 10.1134/S0003683812010115
- Navarro, Q. R., de Oliveira Corrêa, D., Behling, A., Nosedá, M. D., Amano, É., Suzuki, R. M., et al. (2021). Efficient use of biomass and extract of the microalga scenedesmus subspicatus (Scenedesmeceae) in asymbiotic seed germination and seedling development of the orchid cattleya warneri. *J. Appl. Phycol.* 33, 1–19. doi: 10.1007/s10811-021-02442-y
- Navarro-López, E., Cerón-García, M., del, C., López-Rodríguez, M., Acien-Fernández, F. G., and Molina-Grima, E. (2020a). Biostimulants obtained after pilot-scale high-pressure homogenization of scenedesmus sp. grown in pig manure. *Algal Res.* 52, 102123. doi: 10.1016/j.algal.2020.102123
- Navarro-López, E., Ruiz-Nieto, A., Ferreira, A., Gabriel Acien, F., and Gouveia, L. (2020b). Biostimulant potential of scenedesmus obliquus grown in brewery wastewater. *Molecules* 25, 1–16. doi: 10.3390/molecules25030664
- Nayak, M., Swain, D. K., and Sen, R. (2019). Strategic valorization of de-oiled microalgal biomass waste as biofertilizer for sustainable and improved agriculture of rice (*Oryza sativa* L.) crop. *Sci. Total. Environ.* 682, 475–484. doi: 10.1016/j.scitotenv.2019.05.123
- Nilsson, M., Bhattacharya, J., Rai, A. N., and Bergman, B. (2002). Colonization of roots of rice (*Oryza sativa*) by symbiotic nostoc strains. *New Phytol.* 156, 517–525. doi: 10.1046/j.1469-8137.2002.00534.x
- Nisha, R., Kaushik, A., and Kaushik, C. P. (2007). Effect of indigenous cyanobacterial application on structural stability and productivity of an organically poor semi-arid soil. *Geoderma* 138, 49–56. doi: 10.1016/j.geoderma.2006.10.007
- Nisha, R., Kiran, B., Kaushik, A., and Kaushik, C. P. (2018). Bioremediation of salt affected soils using cyanobacteria in terms of physical structure, nutrient status and microbial activity. *Int. J. Environ. Sci. Technol.* 15, 571–580. doi: 10.1007/s13762-017-1419-7
- Osman, M. E. H., El-Sheekh, M. M., El-Naggar, A. H., and Gheda, S. F. (2010). Effect of two species of cyanobacteria as biofertilizers on some metabolic activities, growth, and yield of pea plant. *Biol. Fertil. Soils.* 46, 861–875. doi: 10.1007/s00374-010-0491-7
- Pan, S., Jeevanandam, J., and Danquah, M. K. (2019). “Benefits of algal extracts in sustainable agriculture,”. *Grand Challenges in Algae Biotechnology. Grand Challenges in Biology and Biotechnology*. Eds. A. Hallmann and P. Rampelotto (Cham, Switzerland: Springer), 501–534.
- Pereira, N. S., Ferreira, B. R. R., de Carvalho, E. M., and Damiani, C. R. (2018). Application of chlorella sorokiniana (Chlorophyceae) as supplement and/or an alternative medium for the *in vitro* cultivation of schomburgkia crispa (Orchidaceae). *J. Appl. Phycol.* 30, 2347–2358. doi: 10.1007/s10811-018-1441-2
- Pereira, I., Ortega, R., Barrientos, L., Moya, M., Reyes, G., and Kramm, V. (2009). Development of a biofertilizer based on filamentous nitrogen-fixing cyanobacteria for rice crops in Chile. *J. Appl. Phycol.* 21, 135–144. doi: 10.1007/s10811-008-9342-4
- Plaza, B. M., Gómez-Serrano, C., Acien-Fernández, F. G., and Jimenez-Becker, S. (2018). Effect of microalgae hydrolysate foliar application (*Arthrospira platensis* and scenedesmus sp.) on petunia x hybrida growth. *J. Appl. Phycol.* 30, 2359–2365. doi: 10.1007/s10811-018-1427-0
- Povero, G., Mejia, J. F., Di Tommaso, D., Piaggese, A., and Warrior, P. (2016). A systematic approach to discover and characterize natural plant biostimulants. *Front. Plant Sci.* 7. doi: 10.3389/fpls.2016.00435
- Prasanna, R., Adak, A., Verma, S., Bidyarani, N., Babu, S., Pal, M., et al. (2015a). Cyanobacterial inoculation in rice grown under flooded and SRI modes of cultivation elicits differential effects on plant growth and nutrient dynamics. *Ecol. Eng.* 84, 532–541. doi: 10.1016/j.ecoleng.2015.09.033
- Prasanna, R., Babu, S., Bidyarani, N., Kumar, A., Triveni, S., Monga, D., et al. (2015b). Prospecting cyanobacteria-fortified composts as plant growth promoting and biocontrol agents in cotton. *Exp. Agric.* 51, 42–65. doi: 10.1017/S0014479714000143
- Prasanna, R., Babu, S., Rana, A., Kabi, S. R., Chaudhary, V., Gupta, V., et al. (2013a). Evaluating the establishment and agronomic proficiency of cyanobacterial consortia as organic options in wheat-rice cropping sequence. *Exp. Agric.* 49, 416–434. doi: 10.1017/S001447971200107X
- Prasanna, R., Chaudhary, V., Gupta, V., Babu, S., Kumar, A., Singh, R., et al. (2013b). Cyanobacteria mediated plant growth promotion and bioprotection against fusarium wilt in tomato. *Eur. J. Plant Pathol.* 136, 337–353. doi: 10.1007/s10658-013-0167-x
- Prasanna, R., Gupta, H., Yadav, V. K., Gupta, K., Buddhadeo, R., Gogoi, R., et al. (2020). Prospecting the promise of cyanobacterial formulations developed using soil-less substrates as carriers. *Environ. Technol. Innov.* 18, 100652. doi: 10.1016/j.eti.2020.100652
- Prasanna, R., Hossain, F., Babu, S., Bidyarani, N., Adak, A., Verma, S., et al. (2015c). Prospecting cyanobacterial formulations as plant-growth-promoting agents for maize hybrids. *South Afr. J. Plant Soil* 32, 199–207. doi: 10.1080/02571862.2015.1025444
- Prasanna, R., Kanchan, A., Ramakrishnan, B., Ranjan, K., Venkatachalam, S., Hossain, F., et al. (2016a). Cyanobacteria-based bioinoculants influence growth and yields by modulating the microbial communities favourably in the rhizospheres of maize hybrids. *Eur. J. Soil Biol.* 75, 15–23. doi: 10.1016/j.ejsobi.2016.04.001
- Prasanna, R., Nain, L., Tripathi, R., Gupta, V., Chaudhary, V., Middha, S., et al. (2008). Evaluation of fungicidal activity of extracellular filtrates of cyanobacteria - possible role of hydrolytic enzymes. *J. Basic. Microbiol.* 48, 186–194. doi: 10.1002/jobm.200700199

- Prasanna, R., Pattnaik, S., Sugitha, T. C. K., Nain, L., and Saxena, A. K. (2011). Development of cyanobacterium-based biofilms and their *in vitro* evaluation for agriculturally useful traits. *Folia Microbiol.* 56, 49–58. doi: 10.1007/S12223-011-0013-5
- Prasanna, R., Ramakrishnan, B., Ranjan, K., Venkatachalam, S., Kanchan, A., Solanki, P., et al. (2016b). Microbial inoculants with multifaceted traits suppress *Rhizoctonia* populations and promote plant growth in cotton. *J. Phytopathol.* 164, 1030–1042. doi: 10.1111/jph.12524
- Prasanna, R., Triveni, S., Bidiyaran, N., Babu, S., Yadav, K., Adak, A., et al. (2014). Evaluating the efficacy of cyanobacterial formulations and biofilmed inoculants for leguminous crops. *Arch. Agron. Soil Sci.* 60, 349–366. doi: 10.1080/03650340.2013.792407
- Priya, M., Gurung, N., Mukherjee, K., and Bose, S. (2014). “23 - Microalgae in removal of heavy metal and organic pollutants from soil,” *Microbial Biodegradation and Bioremediation*. ed. S. Das (London, UK: Elsevier), 519–537. doi: 10.1016/B978-0-12-800021-2.00023-6
- Priya, H., Prasanna, R., Ramakrishnan, B., Bidiyaran, N., Babu, S., Thapa, S., et al. (2015). Influence of cyanobacterial inoculation on the culturable microbiome and growth of rice. *Microbiol. Res.* 171, 78–89. doi: 10.1016/j.micres.2014.12.011
- Puglisi, I., La Bella, E., Rovetto, E. I., Lo Piero, A. R., and Baglieri, A. (2020). Biostimulant effect and biochemical response in lettuce seedlings treated with a *scenedesmus quadricauda* extract. *Plants* 9, 123. doi: 10.3390/plants9010123
- Rachidi, F., Benhima, R., Kasmi, Y., Sbabou, L., and El Arroussi, H. (2021). Evaluation of microalgae polysaccharides as biostimulants of tomato plant defense using metabolomics and biochemical approaches. *Sci. Rep.* 11, 930. doi: 10.1038/s41598-020-78820-2
- Ramírez-López, C., Esparza-García, F. J., Ferrera-Cerrato, R., Alarcón, A., and Cañizares-Villanueva, R. O. (2019). Short-term effects of a photosynthetic microbial consortium and nitrogen fertilization on soil chemical properties, growth, and yield of wheat under greenhouse conditions. *J. Appl. Phycol.* 31, 3617–3624. doi: 10.1007/s10811-019-01861-2
- Rana, A., Joshi, M., Prasanna, R., Shivay, Y. S., and Nain, L. (2012). Biofortification of wheat through inoculation of plant growth promoting rhizobacteria and cyanobacteria. *Eur. J. Soil Biol.* 50, 118–126. doi: 10.1016/j.ejsobi.2012.01.005
- Ranglová, K., Lakatos, G. E., Cámara Manóel, J. A., Grivalský, T., Suárez Estrella, F., Acien Fernández, F. G., et al. (2021). Growth, biostimulant and biopesticide activity of the MACC-1 *Chlorella* strain cultivated outdoors in inorganic medium and wastewater. *Algal. Res.* 53, 102136. doi: 10.1016/j.algal.2020.102136
- Rashad, S., El-Hassanin, A. S., Mostafa, S. S.M., and El-Chaghaby, G. A. (2019). Cyanobacteria cultivation using olive milling wastewater for bio-fertilization of celery plan. *Glob. J. Environ. Sci. Manag.* 5, 167–174. doi: 10.22034/gjesm.2019.02.03
- Regulation (EU) 2019/1009 (2019). *REGULATION (EU) 2019/1009 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 5 June 2019 laying down rules on the making available on the market of EU fertilising products and amending regulations (EC) no 1069/2009 and (EC) no 1107/2009 and repealing regula.*
- Renuka, N., Guldhe, A., Prasanna, R., Singh, P., and Bux, F. (2018). Microalgae as multi-functional options in modern agriculture: current trends, prospects and challenges. *Biotechnol. Adv.* 36, 1255–1273. doi: 10.1016/j.biotechadv.2018.04.004
- Renuka, N., Prasanna, R., Sood, A., Ahluwalia, A. S., Bansal, R., Babu, S., et al. (2016). Exploring the efficacy of wastewater-grown microalgal biomass as a biofertilizer for wheat. *Environ. Sci. Pollut. Res.* 23, 6608–6620. doi: 10.1007/s11356-015-5884-6
- Righini, H., Francioso, O., Di Foggia, M., Prodi, A., Quintana, A. M., and Roberti, R. (2021). Tomato seed biopriming with water extracts from *anabaena minutissima*, *ecklonia maxima* and *jania adhaerens* as a new agro-ecological option against *rhizoctonia solani*. *Sci. Hortic. (Amsterdam)*. 281, 109921. doi: 10.1016/J.SCI.2021.109921
- Ronga, D., Biazzi, E., Parati, K., Carminati, D., Carminati, E., and Tava, A. (2019). Microalgal biostimulants and biofertilisers in crop productions. *Agronomy* 9, 192. doi: 10.3390/agronomy9040192
- Saadaoui, I., Sedky, R., Rasheed, R., Bounnit, T., Almahmoud, A., Elshekh, A., et al. (2019). Assessment of the algae-based biofertilizer influence on date palm (*Phoenix dactylifera* L.) cultivation. *J. Appl. Phycol.* 31, 457–463. doi: 10.1007/s10811-018-1539-6
- Saadatnia, H., and Riahi, H. (2009). Cyanobacteria from paddy fields in Iran as a biofertilizer in rice plants. *Plant. Soil Environ.* 55, 207–212. doi: 10.17221/384-pse
- Saber, A. A., Hamed, S. M., Abdel-Rahim, E. F. M., and Cantonati, M. (2018). Insecticidal prospects of algal and cyanobacterial extracts against the cotton leafworm *spodoptera littoralis*. *Vie. Milieu*. 68, 199–212.
- Sahu, D., Priyadarshani, I., and Rath, B. (2012). Cyanobacteria -as potential biofertilizer. *CIBTech. J. Microbiol. Jul.-Sept. Oct.-Dec* 1, 2319–386720. Available at: <http://www.cibtech.org/cjm.htm>. (Accessed May 29, 2023).
- Salvi, L., Niccolai, A., Cataldo, E., Sbraci, S., Paoli, F., Storch, P., et al. (2020). Effects of *arthrospira platensis* extract on physiology and berry traits in *vitis vinifera*. *Plants* 9, 1805. doi: 10.3390/plants9121805
- Schreiber, C., Schiedung, H., Harrison, L., Briese, C., Ackermann, B., Kant, J., et al. (2018). Evaluating potential of green alga *chlorella vulgaris* to accumulate phosphorus and to fertilize nutrient-poor soil substrates for crop plants. *J. Appl. Phycol.* 30, 2827–2836. doi: 10.1007/s10811-018-1390-9
- Shaaban, M. M. (2001a). Green microalgae water extract as foliar feeding to wheat plants. *Pakistan J. Biol. Sci.* 4, 628–632. doi: 10.3923/pjbs.2001.628.632
- Shaaban, M. M. (2001b). Nutritional status and growth of maize plants as affected by green microalgae as soil additives. *J. Biol. Sci.* 1, 475–479. doi: 10.3923/jbs.2001.475.479
- Shamim, A., Mahfooz, S., Hussain, A., and Farooqui, A. (2020). Ability of Al-acclimatized immobilized nostoc muscorum to combat abiotic stress and its potential as a biofertilizer. *J. Pure. Appl. Microbiol.* 14, 1377–1386. doi: 10.22207/JFAM.14.2.35
- Shariatmadari, Z., Riahi, H., Seyed Hashtroudi, M., Ghassempour, A., and Aghashariatmadary, Z. (2013). Plant growth promoting cyanobacteria and their distribution in terrestrial habitats of Iran. *Soil Sci. Plant Nutr.* 59, 535–547. doi: 10.1080/00380768.2013.782253
- Sharma, R., Khokhar, M. K., Jat, R. L., and Khandelwal, S. K. (2012). Role of algae and cyanobacteria in sustainable agriculture system. *Wudpecker. J. Agric. Res.* 1, 381–388.
- Sholkamy, E. N., El-Komy, H. M., and Ali, H. M. (2015). Enhancement of soybean (*Glycine max* L.) growth by bio-fertilizers of *Nostoc muscorum* and *Nostoc rivulare*. *Pakistan J. Bot.* 47, 1199–1204. Available at: [http://www.pakbs.org/pjbot/PDFs/47\(3\)/50.pdf](http://www.pakbs.org/pjbot/PDFs/47(3)/50.pdf). (Accessed May 29, 2023).
- Silambarasan, S., Logeswari, P., Sivaramakrishnan, R., Incharoensakdi, A., Cornejo, P., Kamaraj, B., et al. (2021). Removal of nutrients from domestic wastewater by microalgae coupled to lipid augmentation for biodiesel production and influence of deoiled algal biomass as biofertilizer for solanum lycopersicum cultivation. *Chemosphere* 268, 129323. doi: 10.1016/j.chemosphere.2020.129323
- Silva, D.S.O.e, Rocha, R. H. C., da Nóbrega, J. S., Dias, G. A., de Lima, J. F., and Guedes, W. A. (2017). Post-harvest quality of lettuce cv. Elba in relation to spirulina platensis foliar applications. *Científica* 45, 162–168. doi: 10.15361/1984-5529.2017v45n2p162-168
- Simranjit, K., Kanchan, A., Prasanna, R., Ranjan, K., Ramakrishnan, B., Singh, A. K., et al. (2019). Microbial inoculants as plant growth stimulating and soil nutrient availability enhancing options for cucumber under protected cultivation. *World J. Microbiol. Biotechnol.* 35, 51. doi: 10.1007/s11274-019-2623-z
- Singh, J. S., Kumar, A., Rai, A. N., and Singh, D. P. (2016). Cyanobacteria: a precious bio-resource in agriculture, ecosystem, and environmental sustainability. *Front. Microbiol.* 7. doi: 10.3389/fmicb.2016.00529
- Smith, C., Hill, A. K., and Torrente-Murciano, L. (2020). Current and future role of haber-Bosch ammonia in a carbon-free energy landscape. *Energy Environ. Sci.* 13, 331–344. doi: 10.1039/C9EE02873K
- Soppelsa, S., Kelderer, M., Casera, C., Bassi, M., Robatscher, P., Matteazzi, A., et al. (2019). Foliar applications of biostimulants promote growth, yield and fruit quality of strawberry plants grown under nutrient limitation. *Agronomy* 9, 483. doi: 10.3390/agronomy9090483
- Srivastava, S., Srivastava, S., Bist, V., Awasthi, S., Chauhan, R., Chaudhry, V., et al. (2018). *Chlorella vulgaris* and *pseudomonas putida* interaction modulates phosphate trafficking for reduced arsenic uptake in rice (*Oryza sativa* L.). *J. Hazard. Mater.* 351, 177–187. doi: 10.1016/j.jhazmat.2018.02.039
- Stirk, W. A., Bálint, P., Vambe, M., Lovász, C., Molnár, Z., van Staden, J., et al. (2020). Effect of cell disruption methods on the extraction of bioactive metabolites from microalgal biomass. *J. Biotechnol.* 307, 35–43. doi: 10.1016/j.jbiotec.2019.10.012
- Stirk, W. A., Ördög, V., Van Staden, J., and Jäger, K. (2002). Cytokinin- and auxin-like activity in cyanophyta and microalgae. *J. Appl. Phycol.* 14, 215–221. doi: 10.1023/A:1019928425569
- Supraja, K. V., Behera, B., and Balasubramanian, P. (2020). Efficacy of microalgal extracts as biostimulants through seed treatment and foliar spray for tomato cultivation. *Ind. Crops Prod.* 151, 112453. doi: 10.1016/j.indcrop.2020.112453
- Suresh, A., Soundararajan, S., Elavarasi, S., Lewis Oscar, F., and Thajuddin, N. (2019). Evaluation and characterization of the plant growth promoting potentials of two heterocystous cyanobacteria for improving food grains growth. *Biocatal. Agric. Biotechnol.* 17, 647–652. doi: 10.1016/j.cbac.2019.01.002
- Suresh Kumar, K., Dahms, H. U., Won, E. J., Lee, J. S., and Shin, K. H. (2015). Microalgae – a promising tool for heavy metal remediation. *Ecotoxicol. Environ. Saf.* 113, 329–352. doi: 10.1016/J.ECOENV.2014.12.019
- Swarnalakshmi, K., Prasanna, R., Kumar, A., Pattnaik, S., Chakravarty, K., Shivay, Y. S., et al. (2013). Evaluating the influence of novel cyanobacterial biofilmed biofertilizers on soil fertility and plant nutrition in wheat. *Eur. J. Soil Biol.* 55, 107–116. doi: 10.1016/j.ejsobi.2012.12.008
- Tejada-Ruiz, S., Gonzalez-Lopez, C., Rojas, E., and Jiménez-Becker, S. (2020). Effect of the foliar application of microalgae hydrolysate (*Arthrospira platensis*) and silicon on the growth of *pelargonium hortorum* L.H. bailey under salinity conditions. *Agronomy* 10, 1713. doi: 10.3390/agronomy10111713
- Tripathi, R. D., Dwivedi, S., Shukla, M. K., Mishra, S., Srivastava, S., Singh, R., et al. (2008). Role of blue green algae biofertilizer in ameliorating the nitrogen demand and fly-ash stress to the growth and yield of rice (*Oryza sativa* L.) plants. *Chemosphere* 70, 1919–1929. doi: 10.1016/j.chemosphere.2007.07.038
- Triveni, S., Prasanna, R., Kumar, A., Bidiyaran, N., Singh, R., and Saxena, A. K. (2015). Evaluating the promise of trichoderma and *anabaena* based biofilms as multifunctional agents in macrophomina phaseolina-infected cotton crop. *Biocontrol Sci. Technol.* 25, 656–670. doi: 10.1080/09583157.2015.1006171
- USGS (2021). Map of worldwide croplands. In: *Glob. food secur. anal. data 30 meters proj.* Available at: <https://www.croplands.org/app/map?lat=0.17578&lng=0&zoom=2> (Accessed July 6, 2021).
- Uysal, O., Uysal, F. O., and Ekinci, K. (2015). Evaluation of microalgae as microbial fertilizer. *Eur. J. Sustain. Dev.* 4, 77–82. doi: 10.14207/ejsd.2015.v4n2p77

- Viegas, C., Gouveia, L., and Gonçalves, M. (2021a). Aquaculture wastewater treatment through microalgal biomass potential applications on animal feed, agriculture, and energy. *J. Environ. Manage.* 286, 112187. doi: 10.1016/J.JENVMAN.2021.112187
- Viegas, C., Gouveia, L., and Gonçalves, M. (2021b). Bioremediation of cattle manure using microalgae after pre-treatment with biomass ash. *Bioresour. Technol. Rep.* 14, 100681. doi: 10.1016/J.BITEB.2021.100681
- Viegas, C., Gouveia, L., and Gonçalves, M. (2021c). Evaluation of microalgae as bioremediation agent for poultry effluent and biostimulant for germination. *Environ. Technol. Innov.* 24, 102048. doi: 10.1016/J.ETI.2021.102048
- Win, T. T., Barone, G. D., Secundo, F., and Fu, P. (2018). Algal biofertilizers and plant growth stimulants for sustainable agriculture. *Ind. Biotechnol.* 14, 203–211. doi: 10.1089/ind.2018.0010
- Wuang, S. C., Khin, M. C., Chua, P. Q. D., and Luo, Y. D. (2016). Use of spirulina biomass produced from treatment of aquaculture wastewater as agricultural fertilizers. *Algal. Res.* 15, 59–64. doi: 10.1016/j.algal.2016.02.009
- Xu, Y., Rossi, F., Colica, G., Deng, S., De Philippis, R., and Chen, L. (2013). Use of cyanobacterial polysaccharides to promote shrub performances in desert soils: a potential approach for the restoration of desertified areas. *Biol. Fertil. Soils.* 49, 143–152. doi: 10.1007/s00374-012-0707-0
- Yadavalli, R., and Heggers, G. R. V. N. (2013). Two stage treatment of dairy effluent using immobilized *Chlorella pyrenoidosa*. *J. Environ. Heal. Sci. Eng.* 11, 1–6. doi: 10.1186/2052-336x-11-36
- Zaki, S. S., Belal, E. E. E., and Rady, M. M. (2019). Cyanobacteria and glutathione applications improve productivity, nutrient contents, and antioxidant systems of salt-stressed soybean plant. *Int. Lett. Nat. Sci.* 76, 72–85. doi: 10.18052/www.scipress.com/ilns.76.72
- Zarezadeh, S., Moheimani, N. R., Jenkins, S. N., Hülsen, T., Riahi, H., and Mickan, B. S. (2019). Microalgae and phototrophic purple bacteria for nutrient recovery from agri-industrial effluents: influences on plant growth, rhizosphere bacteria, and putative carbon- and nitrogen-cycling genes. *Front. Plant Sci.* 10. doi: 10.3389/fpls.2019.01193
- Zayadan, B. K., Matorin, D. N., Baimakhanova, G. B., Bolathan, K., Oraz, G. D., and Sadanov, A. K. (2014). Promising microbial consortia for producing biofertilizers for rice fields. *Microbiol. (Russian. Fed.)* 83, 391–397. doi: 10.1134/S0026261714040171
- Zhang, J., Song, X., Wei, H., Zhou, W., Peng, C., and Li, D. (2021). Effect of substituting nitrogen fertilizer with nitrogen-fixing cyanobacteria on yield in a double-rice cropping system in southern China. *J. Appl. Phycol.* 33, 2221–2232. doi: 10.1007/s10811-021-02455-7
- Zilio, M., Pigoli, A., Rizzi, B., Herrera, A., Tambone, F., Geromel, G., et al. (2022). Using highly stabilized digestate and digestate-derived ammonium sulphate to replace synthetic fertilizers: the effects on soil, environment, and crop production. *Sci. Total. Environ.* 815, 152919. doi: 10.1016/J.SCITOTENV.2022.152919